

Review Article

Abundance estimation with DNA metabarcoding – recent advancements for terrestrial arthropods

 Wiebke Sickel¹, Vera Zizka², Alice Scherges², Sarah J. Bourlat², Petra Dieker^{1,3}
¹ Thünen Institute of Biodiversity, Bundesallee 65, 38116 Braunschweig, Germany

² Leibniz Institute for the Analysis of Biodiversity Change, Museum Koenig Bonn, Adenauerallee 127, 53113 Bonn, Germany

³ National Monitoring Centre for Biodiversity, Alte Messe 6, 04103 Leipzig, Germany

 Corresponding author: Wiebke Sickel (wiebke.sickel@thuenen.de)

Abstract

Biodiversity is declining at alarming rates worldwide and large-scale monitoring is urgently needed to understand changes and their drivers. While classical taxonomic identification of species is time and labour intensive, the combination with DNA-based methods could upscale monitoring activities to achieve larger spatial coverage and increased sampling effort. However, challenges remain for DNA-based methods when the number of individuals per species and/or biomass estimates are required. Several methodological advancements exist to improve the potential of DNA metabarcoding for abundance analysis, which however need further evaluation. Here, we discuss laboratory, as well as some bioinformatic adjustments to DNA metabarcoding workflows regarding their potential to achieve species abundance estimation from arthropod community samples. Our review includes pre-laboratory processing methods such as specimen photography, laboratory methods such as the use of spike-in DNA as an internal standard and bioinformatic advancements like correction factors. We conclude that specimen photography coupled with DNA metabarcoding currently promises the greatest potential to achieve estimates of the number of individuals per species and biomass estimates, but that approaches such as spike-ins and correction factors are promising methods to pursue further.

Key words: abundance, biodiversity monitoring, *COI*, insects, metabarcoding, spike-ins



Academic editor: Alfried Vogler

Received: 6 September 2023

Accepted: 1 October 2023

Published: 23 November 2023

Citation: Sickel W, Zizka V, Scherges A, Bourlat SJ, Dieker P (2023) Abundance estimation with DNA metabarcoding – recent advancements for terrestrial arthropods. *Metabarcoding and Metagenomics* 7: e112290. <https://doi.org/10.3897/mbmg.7.112290>

Copyright: © Wiebke Sickel et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Biodiversity is declining at alarming rates worldwide (Díaz et al. 2020). The startling observation of a decline in over 75% flying insect biomass in German nature reserves over 25 years (Hallmann et al. 2017) triggered an earthquake in society and politics and raised awareness of arthropod declines, which have since been further documented (Lister and Garcia 2018; Seibold et al. 2019; Simmons et al. 2019; van Klink et al. 2020). Subsequently, numerous initiatives have been launched or reinforced at global to European and regional scales to assess arthropod diversity and also define guidelines for applied, large-scale biodiversity monitoring schemes (Seibold et al. 2019; Ronquist et al. 2020; Potts et al. 2021). Monitoring programmes are frequently limited in spatial coverage

and sampling effort, since they often rely on morpho-taxonomical analysis for species identifications, which is costly and time-consuming (Yu et al. 2012) and additionally limited by a shortage of taxonomic expertise (Fernandes et al. 2019; Watts et al. 2019; Darby et al. 2020; van Klink et al. 2022a). Thus, in order to meet the increased demand for arthropod diversity assessments, traditional morpho-taxonomy approaches need to be combined with other methods (Pawlowski et al. 2018; Compson et al. 2020).

DNA-based approaches offer a promising alternative to arthropod diversity surveys and monitoring (Porter and Hajibabaei (2018); Zinger et al. (2020); Suppl. material 1). In particular, DNA metabarcoding enables high sample throughput (Elbrecht and Steinke 2018; de Kerdrel et al. 2020), due to automation of laboratory and bioinformatic workflows (Krehenwinkel et al. 2017a; Buchner et al. 2021; Buchner et al. 2023) and offers a time- and cost-effective approach for large-scale biodiversity assessments (Piper et al. 2019; Watts et al. 2019). Molecular methods further have the potential to resolve cryptic species (Sow et al. 2019) and intraspecific genetic diversity (Elbrecht et al. 2018) and open up the possibility to include degraded and non-invasively collected material, e.g. faeces (Andriollo et al. 2019), or plant material in biodiversity surveys, which yields high potential for trophic interaction and food web analysis.

However, implementation in policy-mandated monitoring programmes is still hampered (Blancher et al. 2022; Kelly et al. 2023). Reasons for the limited application include general scepticism among taxonomists, missing expertise and infrastructure within state monitoring agencies, a lack of standardised molecular protocols (Dickie et al. 2018; Pawlowski et al. 2018; Zinger et al. 2019; Compson et al. 2020; Creedy et al. 2021), as well as incomplete reference databases (Watts et al. 2019; van der Heyde et al. 2020; Zenker et al. 2020) and the destruction of specimens for DNA extraction (Zizka et al. 2019), although non-destructive approaches are gaining ground (Castalanelli et al. 2010; Carew et al. 2018; Zenker et al. 2020; Batovska et al. 2021; Kirse et al. 2023). The most important shortcoming concerns the limitation to assess the number of individuals per species and biomass, which is essential in standardised monitoring and ecological analysis, but still remains one of the greatest challenges for high-throughput DNA-based approaches (Compson et al. 2020).

Several factors within the metabarcoding workflow affect extraction of abundance data (Pawlowski et al. 2018; Zinger et al. 2019). Firstly, sample properties such as complexity seem to affect abundance information (Piñol et al. 2019). This complexity refers in particular to variation in biomass across and within species (Elbrecht and Leese 2015; Elbrecht et al. 2017; Braukmann et al. 2019), but also marker gene copy numbers (Krehenwinkel et al. 2017b). Secondly, methodological biases skew abundance and biomass estimations. During DNA extraction, a protocol-dependent taxonomic bias can be introduced due to variations in species size and morphology, causing differences in isolated DNA yields (Krehenwinkel et al. 2017a; Pornon et al. 2017; Matos-Maraví et al. 2019; Iwaszkiewicz-Eggebrecht et al. 2022). Several steps in the metabarcoding laboratory workflow, such as PCR amplification and sequencing, can introduce stochastic processes affecting read counts (Leray and Knowlton 2017; Shirazi et al. 2021). Arguably, the strongest bias is caused by taxon-specific differences in primer binding efficiency (Piñol et al. 2015; Krehenwinkel

et al. 2017a, 2017b). The magnitude of primer bias depends on the number of mismatches between primer and target sequence, especially towards the 3'-end of the primer (Piñol et al. 2019). Apart from primer choice, additional PCR bias can be caused by variable GC content in the target genetic marker (Nichols et al. 2018), amplicon length (Krehenwinkel et al. 2017b) or the occurrence of pseudogenes (Andujar et al. 2021). Thirdly, post-laboratory steps in the bioinformatic processing of sequencing data can skew final read distribution (Frøslev et al. 2017; Alberdi et al. 2018; Darby et al. 2020; Matos-Maraví et al. 2019; Creedy et al. 2021).

A meta-analysis targeting 22 DNA metabarcoding studies revealed a weak relationship between biomass and generated read counts, with a large degree of uncertainty (Lamb et al. 2019). The studies included in Lamb et al. (2019) used different protocols and a wide range of target organisms and sample types, which somewhat hampers overall comparability, but does emphasise that raw read counts are not suitable to infer abundance estimates.

A variety of different approaches have emerged recently that can help improve abundance and biomass estimates from metabarcoding data, including species-specific correction factors applied to read counts, spike-ins, primer optimisation or multi-locus metabarcoding (e.g. Richardson et al. (2015); Krehenwinkel et al. (2017b); Richardson et al. (2019); Darby et al. (2020); Luo et al. (2023); Suppl. material 1). However, these advances have so far not been compared systematically for complex arthropod samples. Furthermore, since taxonomic biases exist in DNA metabarcoding data (Clarke et al. 2014), some form of species-specific correction of DNA metabarcoding would be required to obtain robust estimates of individuals per species. This may not be required to reliably estimate total biomass, but still requires the conversion of read counts to biomass, which is not trivial.

Here, we review potential methods that can improve abundance and biomass estimation in arthropod whole organism community (WOC) samples. Considering the variety of approaches and applications, we aim to formulate general recommendations for DNA metabarcoding workflows in arthropod monitoring. In addition, we explore approaches from metabarcoding studies targeting e.g. aquatic samples that have so far not been applied to terrestrial arthropods and their trophic interactions.

Methods

Collection of relevant literature and assessment of methodological approaches

We performed an online literature search in Google Scholar and EBSCO Discovery Service on 17 January 2022 using the keywords [(quant*) AND (insect) AND (metabarcod*) AND (DNA)] and included only peer reviewed publications in English. Although the search term specifically targeted insects, we use the more general term “terrestrial arthropods” throughout the text. Additionally, some publications were added to the list based on the authors' expertise.

We included studies that applied DNA metabarcoding to terrestrial arthropods as target organisms and/or in relation to their trophic interactions within ecosystems (e.g. pollination and food web studies), as these topics are strongly

connected and play an important role in monitoring schemes (e.g. ecosystem services of pollination or natural pest control). With these criteria, WOC and tissue samples were included covering also pollen, gut contents and faeces as well eDNA metabarcoding approaches, such as extraction from soil and sample fixative. We excluded studies that applied individual-based DNA barcoding and next generation sequencing (NGS) barcoding, PCR-free approaches as well as long-read sequencing methodologies, as we wanted to focus on metabarcoding specifically. PCR-free approaches are, however, briefly discussed in an outlook section.

Based on 113 publications matching our search criteria (Suppl. material 2), we extracted information on article type, study type, sample type, species group, methods and parameters (Table 1, Suppl. material 3). We examined these methods regarding their applicability to study types (species richness assessments, pollen analysis, food web studies) and to sample types (WOC samples, pollen, eDNA and gut contents/faeces). The overall suitability was assessed based on whether certain abundance metrics (number of individuals per species, relative abundance) as well as biomass estimation were achievable, whilst also considering the extent of additional equipment, cost and labour (Suppl. material 4). These considerations are based on the available literature.

Abbreviations:

ASV	amplicon sequence variant
ddPCR	digital droplet PCR
eDNA	environmental DNA
FOO / POO	frequency of occurrence / percent of occurrence
NGS / HTS	next generation sequencing / high-throughput sequencing
qPCR	quantitative PCR
RRA	relative read abundance
UMI	unique molecular identifier
WOC samples	whole organism community samples

Results

Description and assessment of methods

Reviewing the literature, we identified three main methods to estimate species abundance with metabarcoding (Table 1): (i) semi-quantitative metrics (Fig. 1), (ii) approaches that can potentially reduce read abundance biases (Fig. 2), and (iii) the combination of DNA (meta-)barcoding with other methodological approaches, which we present in more detail in the following sections (also see Suppl. material 1). This review focuses on studies including developments associated with the laboratory workflow. For a critical assessment of missing standards in bioinformatics we refer to Creedy et al. (2021). Since many metabarcoding studies refer to relative abundances, whilst monitoring aims to determine counts of individuals per species, we make a clear distinction of these terms throughout this manuscript by referring to “number of individuals per species” (absolute number of individuals belonging to the same

Table 1. Overview of methodological approaches discussed in this publication.

Category	Approach	Sample types	Quantitative?
Semi-quantitative metrics	FOO/POO	all	semi-quantitative
	RRA	all	semi-quantitative
	rarefaction	all	semi-quantitative
	transformation	all	semi-quantitative
Reducing read abundance biases	correction factors via algorithm	WOC samples	yes; virtual specimen counts
	correction factors via mock communities	all	yes
	spike-ins	all	yes; relative abundance
	primer optimisation	all	no
	multi-locus metabarcoding	all	yes
Combination of methods	general	all; depending on approach	yes; depending on approach
	photography and body measurements of single specimens	WOC samples	yes; number of individuals per species, biomass

species) and “relative abundance” (proportion of a species within a sample), but use the more general term “abundance” as a summary term. “Biomass” (weight of individuals belonging to the same species) is mentioned separately, where appropriate.

Semi-quantitative metrics

DNA metabarcoding is comprehensively used to assess presence/absence from complex sample mixtures. Whilst this can be informative for some ecological assessments, including biodiversity measures (e.g. alpha diversity), interaction analyses (e.g. multi-trophic networks, food web structures, plant-pollinator interactions) require some form of (semi-)quantitative data. There are different approaches to conduct semi-quantitative analysis of DNA metabarcoding data (Fig. 1A). In diet analyses, frequency or percentage of occurrence (FOO/POO; Fig. 1B) are often applied (Deagle et al. (2019), but see Cuff et al. (2022)). In bipartite networks, link strength (Fig. 1C) is a meaningful quantitative metric for plant-pollinator or prey-predator networks (Cuff et al. 2022; Thomsen and Sigsgaard 2019). Alternatively, relative read abundance (RRA) summarised over biological replicates is often used (Fig. 1D), especially for pollen samples (Kratschmer et al. 2019; Wilson et al. 2021). Various studies have applied read counts, RRA as well as derived metrics, such as log- or rank-transformed or rarefied read abundance to assess community composition for different sample types (pollen, faeces, gut and WOC samples; Hope et al. (2014); Hawkins et al. (2015); Richardson et al. (2015); Krehenwinkel et al. (2018); Macías-Hernández et al. (2018); Marquina et al. (2019)). The use of any of the above-mentioned metrics is straightforward, but in most cases they are unreliable for quantitative inferences due to the various factors affecting read counts (Pawlowski et al. 2018; Zinger et al. 2019) and thus they are uninformative to estimate the number of individuals per species or biomass.




Figure 1. Semi-quantitative metrics. **A** ASV table as the outcome of a DNA metabarcoding experiment, rows are samples, columns are ASVs, numbers are raw read counts. From the ASV table, semi-quantitative metrics can be derived, e.g. frequency and percentage of occurrence, bipartite networks and relative read abundance. **B** Frequency and percentage of occurrence derived from ASV table, frequency of occurrence simplifies the ASV table into presence/absence data, indicated by presence or absence of a rectangle (left), when summarising this over all samples, percentage of occurrence can be an informative metric for abundance in a system (right). **C** Bipartite networks derived from the ASV table, samples and ASVs are nodes, edges indicate presence/absence of the ASVs per sample (left), when summarising this over all samples, link strength can be an informative metric for abundance in a system (right). **D** Relative read abundance derived from ASV table, relative read abundance for individual samples is determined by dividing raw read counts of individual ASVs by total read count per sample (left), when summarizing this over all samples, mean relative read abundance can be an informative metric for abundance in a system (right); abbreviations: S – Sample, ASV – Amplicon sequence variant, RRA – relative read abundance; ASVs are colour coded and refer to ASVs from (A), artwork: Alice Scherges.


Approaches that reduce read bias

Correction factors

Mock community experiments have shown a positive correlation of read counts per species with genomic template DNA concentration in pollen and WOC samples (Baksay et al. 2020; Gueuning et al. 2019), while other studies revealed a PCR bias introduced through taxon- and marker-specific primer efficiency (Bell et al. 2019; Braukmann et al. 2019; Darby et al. 2020; Krehenwinkel et al. 2017b). Since these biases are strongly affected by primer binding efficiencies (Piñol et al. 2019), they are assumed to be predictable (Krehenwinkel et al. 2017b). Thus, correcting read counts using species-specific correction factors can improve metabarcoding-derived abundance estimates (Krehenwinkel et al. (2017b); Darby et al. (2020); Fig. 2A, B). Such correction factors can be obtained using mock communities (Krehenwinkel et al. (2017b), Fig. 2A) or iterative “guess-and-test” algorithms (see further down, Darby et al. (2013); Darby et al. (2020); Table 1, Fig. 2B, Suppl. material 1). In order to derive correction factors using mock communities, artificial community samples of defined composition are processed alongside unknown samples. However, the derived correction factors can only be applied to species that are present both in environmental and artificial community samples, which is a strong limitation for hyperdiverse WOC arthropod samples such as Malaise trap catches that contain many unknown taxa.

It may be possible to extend correction factors to closely-related taxa based on phylogenetic relatedness, whereby similar skews of read counts are assumed. In microbial analyses (Goberna and Verdu 2016), such an approach has been used to infer functional traits. In beetles, a recent paper has shown a significant correlation between species biomass and high-throughput sequencing (HTS) read abundance for *16S*, but not for *COI*, which had more primer-template mismatches. In addition, strong phylogenetic signals in primer-template mismatches were identified and models incorporating the effects of mismatch type or number improved species biomass estimation from HTS read abundance for *COI* (Liu et al. 2023). To obtain correction factors, mock communities have so far only been used in combination with WOC arthropod samples (Krehenwinkel et al. 2017b), but this approach could also be transferred to pollen samples, as processing mock communities alongside such samples is common (Bell et al. 2019; Baksay et al. 2020; Swenson and Gemeinholzer 2021). Species-specific correction factors obtained from mock community samples are helpful to reduce read abundance biases; however, some sources of bias still exist, e.g. related to the evenness of a community sample (Piñol et al. 2019), copy number variations of the target gene (Krehenwinkel et al. 2017b) or differences in DNA quality between specimens used for mock community samples versus field-collected samples (Krehenwinkel et al. 2018). Furthermore, it should be noted that correction-factors obtained via mock communities are expected to fluctuate in response to changes in PCR cycle numbers across metabarcoding workflows (Yang et al. 2020; Martoni et al. 2022).

Correction factors can also be calculated using an iterative algorithm which mitigates data skews due to copy number variations of the target gene (Darby et al. (2020, 2013); Fig. 2B, Suppl. material 1). This requires a reference dataset



© Alice Scherges

Figure 2. Reducing read abundance biases. **A** Processing mock communities (bottle) with defined composition allows determining taxon-specific correction factors, which can be applied to correct relative read abundance of samples with unknown composition, indicated by a red line. Correction factors can only be determined for taxa included in the mock community. **B** Correction factors can be determined using iterative algorithms and a guess-and-test approach based on a morphological reference data set (not shown). The correction factors can be applied to correct relative read abundance of samples with unknown composition, indicated by a red line. Correction factors can only be determined for samples that show a good agreement in terms of taxa detected between the reference and the DNA metabarcoding data set. **C** Adding spike-ins, e.g. a defined amount of genomic DNA, to all samples and co-amplifying and co-sequencing the reference material allows correcting raw read counts by simply dividing read counts assigned to taxa (blue and brown bars) by read counts assigned to the spike-in (red bars); abbreviations: RRA – relative read abundance, S – sample, artwork: Alice Scherges.

with specimen counts obtained from morphological identifications. The algorithm itself then starts with randomly generated correction factors for each species to compute predicted specimen counts from metabarcoding data. The predicted counts are compared to the reference data set and correction factors are iteratively adjusted until predicted and actual counts converge (Darby et al. 2013).

The algorithm can only be applied to samples with high concordance between morphological and DNA-based taxonomies, but it is a promising approach, as the predicted numbers of individuals per species were highly correlated with actual count data (Darby et al. 2013; Darby et al. 2020). It requires high-quality material and specimens to be identified morphologically (Darby et al. 2013; Darby et al. 2020) and thus can only be used for WOC and tissue samples (see Suppl. material 1). Time and cost of the overall analysis increases, as a reference set of morphologically identified species is required, but this could be worth it in the case of repeated monitoring of sites with known species composition, or for the monitoring of known arthropod pests.

Spike-ins

Spike-ins (Fig. 2C) may also be referred to as internal standards (ISDs; Harrison et al. (2021)). Here, a defined amount of reference material DNA is added to each sample, which allows read count correction and thereby improves abundance and biomass estimation (Luo et al. 2023). The reference DNA can be added as tissue (Darby et al. 2020), genomic DNA, pre-amplified DNA (Ji et al. 2020), plasmids (Luo et al. 2023) or synthetic DNA (Palmer et al. 2018), should include primer binding sites and needs to be added to the reference database (Tkacz et al. 2018; Luo et al. 2023). Spike-ins are added to the samples in a standardised manner, e.g. a defined amount of reference DNA (ng) per defined volume of lysis buffer (μ l; Ji et al. (2020); Luo et al. (2023)). It is recommended to add the spike-in after tissue lysis but prior to DNA extraction (Ji et al. 2020; Luo et al. 2023), so that it is co-extracted, co-amplified and co-sequenced along with the sample DNA and therefore underlies the same methodological biases. Since all samples receive the same amount of spike-in, they should theoretically return the same spike-in read counts. However, sample complexity affects read numbers (Piñol et al. 2019) and thus different samples will return different read numbers for the spike-in (Luo et al. 2023). Read correction can be achieved by dividing the number of reads assigned to amplicon sequence variants (ASVs) by the number of reads assigned to the spike-in, resulting in significant improvement in within-species abundance across samples (Ji et al. 2020; Luo et al. 2023).

The use of spike-ins is not restricted by sample type, but comes with a low increase in effort and costs, because the spiking of samples is an additional, albeit minimal, step in the laboratory workflow, which has to then be integrated in the bioinformatic workflow. It should be noted that spike-in correction does not correct for biases across species within samples (Luo et al. 2023). The underlying reasons have, to our knowledge, not been systematically addressed, but may very well relate back to sample complexity and primer binding efficiencies (Piñol et al. 2019). It has been proposed that species-specific correction factors obtained from mock communities (see previous section), unique molecular identifiers (UMIs, see outlook section) as well as the application of less biased primers can be used

to correct for within-sample across species biases (Ji et al. 2020; Luo et al. 2023). Spike-in correction is a straightforward and powerful approach with high potential to improve abundance and biomass estimations via DNA metabarcoding.

Primer optimisation

A variety of studies have shown that primer design is an essential part determining the success of DNA metabarcoding studies, both in terms of taxon recovery and read abundance biases (Esnaola et al. 2018; Jusino et al. 2018; Lafage et al. 2019; Pedro et al. 2020). Primers used in DNA metabarcoding need to be universal and the fragment length needs to be suitable for the sequencing platform of choice, whilst allowing for species-level identification (Meusnier et al. 2008). Over and under amplification of different lineages of arthropods (Krehenwinkel et al. 2017b; Darby et al. 2020) as well as certain plant species dominating pollen samples (Bell et al. 2019; Baksay et al. 2020) have been reported and should be minimised as much as possible. Thus, primer design, including primer validation and evaluation, is a painstaking task, which needs to be continuously pursued using complex mock communities to ensure the best choice of primer for arthropod metabarcoding studies (Elbrecht et al. 2019).

Multi-locus metabarcoding

Different genetic markers suffer from different taxonomic biases and thus some studies employ several different loci for the same organismal group, which is referred to as multi-marker (Adamowicz et al. 2019) or multi-locus metabarcoding (Batovska et al. 2021). Multi-locus metabarcoding has been applied to WOC and tissue samples (Marquina et al. 2019; Giebner et al. 2020), pollen (Richardson et al. 2015; Bell et al. 2019; Richardson et al. 2019), faecal samples and gut contents (Swift et al. 2018; Krehenwinkel et al. 2019; Gil et al. 2020) as well as soil and even eDNA samples (Ritter et al. 2019; Thomsen and Sigsgaard 2019).

Locus-specific biases can be mitigated by using rank order abundance or median-based proportional abundance summarised over all loci, as has been demonstrated in pollen DNA metabarcoding (Richardson et al. 2015; Richardson et al. 2019). The locus-specific PCRs are often performed separately (Richardson et al. 2015; Swift et al. 2018; Richardson et al. 2019; Baksay et al. 2020; Darby et al. 2020), which increases time and cost for sample processing. Multi-locus metabarcoding can be performed in multiplexed reactions (de Kerdrel et al. 2020; Batovska et al. 2021) to improve time and cost efficiency. However, this may introduce additional read abundance skews, possibly due to PCR competition between loci (Batovska et al. 2021). During analysis, data from different markers need to be analysed separately (Thomsen and Sigsgaard 2019), which increases time for analysis. It should be emphasised that different markers usually yield discordant taxon lists (Alberdi et al. 2018; da Silva et al. 2019), e.g. because of incomplete reference databases for markers other than *COI* (Andujar et al. 2018), but may also be attributed to differences in PCR efficiency. Such discordant taxa lists allow a broader taxon coverage, but it also means that data from different markers are complementary (Kirse et al. 2021), complicating data analysis. In the case of discordant taxa lists, abundance estimates (e.g. rank-based) can only be determined for taxa identified by more than one marker (Richardson et al. 2015; Richardson et al. 2019).

For pollen samples, as no single universal plant barcode exists (CBOL Plant Working Group 2009; Bell et al. 2016), multi-locus metabarcoding was commonly adopted especially in early studies (Kraaijeveld et al. 2015; Richardson et al. 2015; Pornon et al. 2016). Consequently, pollen DNA metabarcoding workflows include both ribosomal (e.g. *ITS2*) and chloroplast markers (e.g. *trnL*, *rbcL*, *matK*) (Richardson et al. 2015; Milla et al. 2021; Swenson and Gemeinholzer 2021; Arstingstall et al. 2023). The latter generally perform well concerning PCR amplification and taxonomic resolution, despite concerns regarding little chloroplast DNA in pollen grains (Hawkins et al. 2015; Kraaijeveld et al. 2015; Bell et al. 2016). There even exists some evidence that chloroplast markers are more suitable for assessing relative abundances than ribosomal markers (Richardson et al. 2019; Baksay et al. 2020), possibly due to lower GC contents (Baksay et al. 2020), and these differences need to be carefully considered. However, more recently, pollen DNA metabarcoding studies may revert to single-locus metabarcoding, focusing on *ITS2* (Leidenfrost et al. 2020), due to incomplete reference databases for and sub-optimal taxonomic resolution of chloroplast markers (Richardson et al. 2019; Kolter and Gemeinholzer 2021).

Combining DNA metabarcoding with other methods

Some studies combine DNA metabarcoding with other methodologies. Thereby, DNA metabarcoding may be used to obtain a comprehensive species list of the detected taxa, whilst abundance estimates (e.g. number of individuals per species, DNA copy number) and/or biomass estimates are obtained with another methodology. One common example is the complementary morphological analysis of gut content remains, pollen grains or arthropod specimens (Keller et al. 2015; Darby et al. 2020; Gil et al. 2020). Other examples are weighing WOC samples (Hausmann et al. 2020), using flow cytometry of pollen (Baksay et al. 2020) or other forms of PCR (Schneider et al. 2016; Tedersoo et al. 2019). The choice of additional methodology determines the sample types that can be used, for example, combining metabarcoding with quantitative PCR (qPCR; Schneider et al. (2016)) or digital droplet PCR (ddPCR; Tedersoo et al. (2019)) can be performed on all sample types. For other methodologies, for example weighing, WOC samples are required (Hausmann et al. 2020). All these data sources are complementary and can provide different kinds of information, e.g. total biomass of WOC samples obtained from weighing, supplemented by species-level presence/absence data provided by metabarcoding (Hausmann et al. 2020).

One noteworthy approach of method combination is the photographic documentation of specimens from WOC samples before analysing them with DNA metabarcoding. This combined approach enables individual counts, body size measurements and thereby biomass estimation (Gueuning et al. 2019). As specimens are handled individually (Wühl et al. 2022), the use of body parts for DNA extraction, instead of full specimens, is furthermore facilitated (Gueuning et al. 2019; Darby et al. 2020), keeping voucher specimens mostly intact. Specimen photography further allows documentation of specimens for future reference as well as incorporating a pre-sorting strategy (Elbrecht et al. 2020). Whilst handling of individual specimens is exceptionally time- and labour-intensive, automated solutions can improve time-efficiency (Ärje et al. 2020; Wühl et al. 2022). In combination with machine learning approaches, the

automated screening of high-resolution pictures of arthropod WOC samples for abundance estimation is emerging and would facilitate large-scale assessments, e.g. for monitoring schemes (Høye et al. 2021). While these approaches are still in development, the vision of completely automated protocols, incorporating image recognition before molecular sample processing, exists (Høye et al. 2021; Besson et al. 2022; van Klink et al. 2022b; Wühl et al. 2022). However, exactly when highly accurate image recognition to species level for all arthropod species in a WOC sample might be possible and thereby circumvent the need for metabarcoding altogether, is difficult to assess.

Discussion

General conclusions and recommendations for arthropod monitoring and related questions

The available literature has revealed that the majority of (terrestrial) arthropod DNA metabarcoding studies do not sufficiently address the matter of estimating the number of individuals per species and/or biomass (Suppl. material 2). In terms of pollen analysis, research exists that discusses abundance estimation via DNA metabarcoding, but with inconsistent results (Keller et al. 2015; Kraaijeveld et al. 2015; Richardson et al. 2015; Bell et al. 2019; Richardson et al. 2019; Baksay et al. 2020). In contrast to this, DNA metabarcoding has received considerably more attention in the aquatic sector in recent years and advancements exist both for WOC samples of macrozoobenthos and eDNA (Elbrecht and Leese 2015; Elbrecht et al. 2017; Beentjes et al. 2019; Hoshino et al. 2021). Existing policies, like the EU Water Framework Directive (WFD; Directive 2000/60/EC) and the Marine Strategy Framework Directive (MSFD; Directive 2008/56/EC), legally require routine monitoring of aquatic environments. As a consequence, standards for sampling, processing and reporting already exist (Haase et al. (2004), but see Birk et al. (2012)), as well as DNA-based indicators (Aylagas et al. 2014). Especially the DNAqua-Net COST Action (Leese et al. 2016; Leese et al. 2018) has published many advancements regarding the suitability and integration of (e)DNA metabarcoding in bio-monitoring (Buchner et al. 2019; Pawlowski et al. 2018), as well as resources to facilitate standardisation and quality control for DNA-based monitoring (Bruce et al. (2021); Bruce and Keskin (2021); Vasselon et al. (2021); Leese et al. (2023), DNAqua-Hub, <https://dnaquahub.eu/>; accessed 24 May 2022). This work has a high potential to be transferred into terrestrial arthropod monitoring and demonstrates that DNA metabarcoding can indeed be standardised for monitoring purposes (Leese et al. 2023), which is far more challenging for morphological species identification. However, this transfer could be hampered by the lack of data on diversity and distributions of the hyperdiverse arthropods, although such studies are now emerging (Buchner et al. 2023; Srivathsan et al. 2023).


Additionally, the collected literature focused on approaches that apply to the sample processing stage of metabarcoding workflows. The effect of bioinformatics and data analysis strategies on abundance and biomass estimations is strongly underrepresented (Suppl. material 2). A variety of non-harmonised bioinformatic tools and pipelines exists (Creedy et al. 2021), but a more detailed discussion on the bioinformatics and data analysis side of this topic is outside the scope of this review. However, future research needs to address this.

As expected, there is a variety of adjustments attempting to improve abundance and biomass estimation via DNA metabarcoding (Suppl. material 2). It remains, however, difficult to find a “one-size-fits-all” approach to assessing individual counts and biomass from DNA metabarcoding, partly because different approaches are applicable only to certain sample types or because recent advancements still do not translate to individual counts and/or biomass estimates.

Overall suitability of DNA metabarcoding approaches to estimate the number of individuals per species and biomass in terrestrial arthropod monitoring

Currently, the most promising approach is to combine DNA metabarcoding with specimen photography, which would ideally be automated (Ärje et al. 2020; Wühl et al. 2022). In addition, promising avenues such as correction factors and spike-ins should be further developed (Darby et al. 2013; Krehenwinkel et al. 2017b; Darby et al. 2020; Ji et al. 2020; Luo et al. 2023). Specimen photography coupled with automatic image recognition facilitates body size measurements to achieve biomass estimates as well as the number of individuals per species. Combining the approaches of Gueuning et al. (2019), Darby et al. (2020) and de Kerdrel et al. (2020) seems especially promising, as recombining specimens to “pseudo-community” samples allows cost-efficient mixed-species DNA (meta-)barcoding. We would like to point out that this strategy is not the same as NGS barcoding (Wang et al. 2018; Srivathsan et al. 2021), since individual specimens or parts of them are combined to mixed-species samples (Gueuning et al. 2019; de Kerdrel et al. 2020). Thus, samples are processed following a metabarcoding workflow, but obtained barcodes can be traced back to specimens (de Kerdrel et al. 2020). We argue that despite the increase in processing time and associated costs, (automated) specimen photography is a simple and effective way to achieve considerable improvement in taxon recovery, as well as estimates regarding the number of individuals per species and biomass (Fig. 3). This approach is limited to WOC samples, although a similar approach may potentially be applied to pollen samples, for example by flow cytometry (Baksay et al. 2020; Dunker et al. 2020). Theoretically, these approaches could be combined to achieve count and biomass data, although the above-mentioned studies did not comment on this potential.

Regardless of application or sample type, general recommendations for every metabarcoding workflow are to use appropriate positive controls, i.e. mock communities (Ji et al. 2020), as well as negative controls, biological and technical replicates (Alberdi et al. 2018; Elbrecht and Steinke 2018; Liu et al. 2019; Zinger et al. 2019; Yang et al. 2020) and consider multi-locus metabarcoding. Each of these steps can improve taxon detection and the correlation between relative read abundances and input DNA mass (Richardson et al. 2019; Ritter et al. 2019; Thomsen and Sigsgaard 2019; Ji et al. 2020). Associated increases in costs and labour are justified by the improvement in the generated data, although budget limitations may deem technical replicates unfeasible. With optimised metabarcoding and bioinformatic workflows, more robust relative abundance and biomass estimates are thus potentially achievable in the foreseeable future. However, the number of individuals per species cannot be determined, as other sources of bias still exist. We therefore recommend considering additional approaches discussed further down.



© Alice Scherges

Figure 3. Recommended workflow for biodiversity assessments with bulk samples and DNA metabarcoding that obtains count and biomass data with species level taxonomic identifications. **A** Specimens from a bulk sample (bottle) are first processed individually. **B** Processing includes specimen photography (camera), specimen counts (abacus), body size measurements (caliper) and biomass estimation (scales). Ideally, this is done automatically (green robot icon) and involves automatic image recognition to achieve preliminary taxa identifications on broad taxonomic scales. **C** Specimens are then re-combined to a community sample, a spike-in is added and DNA is extracted (microcentrifuge tube). **D** DNA metabarcoding delivers species level identifications and raw read counts (ASV table), which are corrected via the spike-in. **E** Image data is combined to a taxon list containing count, size and biomass data (taxon list). **F** Image data and DNA metabarcoding data are combined using machine learning approaches (data assembly, orange robot icon) to obtain a data set that contains information on species level identities, along with count data and biomass estimates (taxa bubbles), abbreviations: ASV – amplicon sequence variant, artwork: Alice Scherges.

For eDNA, obtaining count data is extremely difficult. Since eDNA dynamics (Barnes and Turner 2016; Compson et al. 2020) are affected by various uncontrollable factors prior to sampling, analysis of abundance information is further impeded. Thus, presence/absence and derived frequency or percentage of occurrence (FOO/POO) data from replicates currently seem to be the best option, although promising approaches exist that will move towards more informative data obtainable from eDNA. For example, combining species detections with information about the cellular and molecular state of eDNA (e.g. intra- versus extra-cellular eDNA, genetic region, fragment size) is expected to improve the abundance estimation, as demonstrated in water samples (Jo et al. 2021). Other options for eDNA-based monitoring are: an overall experimental design and sampling strategy that allows indirect counts, developing and applying novel metrics (e.g. the “eDNA index”; Kelly et al. (2019)) or coupling presence/absence data with site-occupancy models (van Strien et al. 2010; van Strien et al. 2013). We argue that eDNA approaches are worth considering for arthropod monitoring, as they are non-invasive (Andriollo et al. 2019; Thomsen and Sigsgaard 2019; Pumkæo et al. 2021; Roger et al. 2022), which is especially important for protected and endangered species.

Additionally, (e)DNA-based analyses open up new avenues that move away from the traditional estimation of numbers of individuals per species or biomass. One such avenue to pursue further is more sensitive detection rates of parasitism and invasive species (Sow et al. 2019; Young et al. 2021). Thus, (e)DNA metabarcoding deserves to be incorporated in such schemes at least as a complementary approach to morpho-taxonomy.

Outlook: Further molecular approaches for the estimation of species abundances

In the following, we explore selected approaches from the wider literature that were not within the scope of the present review. However, there is high potential for the implementation in monitoring programmes in the future. Novel data analysis pipelines are constantly being developed and some focus on integrating uncertainties associated with the dynamics of DNA in the environment (Barnes and Turner 2016; Compson et al. 2020). One such example, a tracer model, has successfully been applied to estimate the abundance of target fish species (Fukaya et al. 2020). Another example, an “eDNA index”, which is a double-transformation of read-counts, holds potential to assess abundance trends across time and space (Kelly et al. 2019). Additionally, species occupancy models can detect false negatives (Compson et al. 2020) and Bayesian hierarchical models can integrate primer choice and other parameters of the metabarcoding workflow (Doi et al. 2019; Compson et al. 2020), which would allow correcting read count-derived abundance estimates. Lastly, the application of half-life corrections and prey DNA decay rates allow the inference of relative frequencies and biomass of prey items based on metabarcoding and shotgun sequencing data (Uiterwaal and DeLong 2020; Paula et al. 2023).

When grouping sequencing reads as ASVs instead of molecular operational taxonomic units, DNA metabarcoding can potentially deliver conservative abundance estimates in the sense of “minimum census estimates”, similar to those obtained from non-invasive sampling of hair and faeces (Frantz et

al. 2004; Miotto et al. 2007). In this case, the evolutionary rate of the chosen marker would have to be considered (Wang et al. 2016), as it may affect the recovery of ASVs per species and consequently the obtained minimum census estimates. Furthermore, ASV-based analyses facilitate the analysis of intraspecific diversity (Elbrecht et al. 2018; Arribas et al. 2021; Shum and Palumbi 2021; Weitemier et al. 2021), which is greatly underappreciated in arthropod monitoring schemes. Another promising approach is to further refine the qSeq protocol (Hoshino and Inagaki 2017; Hoshino et al. 2021) and similar workflows employing unique molecular identifiers (UMIs; Luo et al. (2023)). Here, a single-primer extension is included in the workflow before performing PCR. During this step, each DNA fragment is labelled with a random tag and the number of random tags per ASV can be used to accurately infer starting copy numbers of each recovered sequence in the original sample. This allows simultaneous species identification and inference of relative abundances from eDNA and WOC samples (Hoshino and Inagaki 2017; Hoshino et al. 2021; Luo et al. 2023). Unique molecular identifiers have also been applied in detecting rare allele variants and mutations and have been reported as being especially useful for read error corrections (Jabara et al. 2011; Kinde et al. 2011; Kivioja et al. 2012; Fields et al. 2020).

There is an urgent need to shift away from a purely morpho-taxonomic approach and related indicators for long-term arthropod monitoring, towards an integrative framework, in which morphological and molecular biological methodologies are applied in parallel. This requires the development and implementation of novel proxies and indicators to indirectly assess species abundance based on genetic data. One possible approach is to apply Hill numbers to DNA-based and morpho-taxonomic assessments alike, as this improves comparability and they can even be applied to (phylo-)genetic data (Alberdi and Gilbert 2019). Additionally, the amount of genomic DNA per taxon can reliably be assessed by combining metabarcoding with qPCR or ddPCR, although these usually focus on specific target species (Schneider et al. 2016; Tedersoo et al. 2019), but also via the use of UMIs (see above).

PCR-free methods represent a further alternative (Garrido-Sanz et al. 2020; Ji et al. 2020; Cordier et al. 2021). The advantage of these approaches is that no amplification step is conducted and therefore, the complete mitochondrial or nuclear DNA is sequenced and analysed. As PCR amplification is omitted, mito- and metagenomic approaches are associated with more reliable abundance estimations. Ideally, whole mitochondrial or nuclear reference genomes of target taxa exist in order to assign generated reads to the species of origin (Schmidt et al. 2022; Theissinger et al. 2023). However, the absence of reference genomes for non-model organisms poses a barrier to the application of PCR-free approaches (Formenti et al. 2022; Lewin et al. 2022) and thus *de novo* assemblies (Meng et al. 2019) may be a suitable alternative, especially for mitogenomics. The general suitability of mito- or metagenomics for large-scale arthropod monitoring, however, remains limited due to higher costs, computing power and data storage requirements. Further, bias introduced through extraction and variable gene copy number still exist in those approaches. Nonetheless, genomic approaches are valuable and informative for biodiversity conservation purposes, as they enable more detailed analyses of intra-specific diversity and population structure than DNA (meta-)barcoding allows (Theissinger et al. 2023).

Concluding remarks

Even though there are many details to consider when applying DNA metabarcoding to arthropod monitoring, pollen and food web analyses, we were able to make some general recommendations. Generally, DNA metabarcoding should always be optimised for maximum taxon recovery and minimal amplification biases. The processing of adequate positive and negative controls is essential. Incorporating appropriate biological and technical replicates reduces the impact of certain methodological biases.

DNA metabarcoding as a rapid tool to obtain species occurrences is a very promising method for large-scale monitoring activities, especially when abundance estimates are not required. When combining DNA metabarcoding with specimen photography and body size measurements, the number of individuals per species and biomass can also be assessed.

Going forward, creating new DNA-based metrics to report (relative) abundances based on genetic units rather than processing individual specimens offers new innovations addressing the most central questions in arthropod monitoring, as these rarely require absolute measures of abundance. Detecting and assessing trends in monitoring relates more to within- and between-sample comparisons taken across spatial and temporal scales, which can be achieved with metabarcoding. Additionally, DNA metabarcoding facilitates the assessment of ecosystem services in a time- and cost-efficient manner, via processing pollen and food web analyses.

There are still many challenges to face until metabarcoding data can deliver robust abundance and biomass estimations. Currently, sorting and individual handling of specimens from WOC samples is unavoidable to obtain such data. However, it is important to apply both classical morpho-taxonomy and molecular biological approaches in parallel, which will allow the management and analysis of the large amounts of data generated by monitoring programmes in a timely and cost-effective manner. Thus, despite its limitations, DNA metabarcoding can and should be incorporated as an additional tool in routine arthropod monitoring to increase sample sizes and cover a broader range of taxonomic groups.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The presented study is part of the joint project "*Monitoring of biodiversity in agricultural landscapes*" (MonViA) that has been funded by the German Federal Ministry of Food and Agriculture.

Author contributions

WS and PD devised the study. WS performed the literature review and drafted the first version of the manuscript. WS, PD, VZ and SJB were substantially involved in subsequent drafts. AS created the figures. All authors agreed to the final version of the manuscript.

Author ORCIDs

Wiebke Sickel  <https://orcid.org/0000-0002-0038-1478>

Vera Zizka  <https://orcid.org/0000-0001-8486-8883>

Alice Scherges  <https://orcid.org/0009-0002-0824-7991>

Sarah J. Bourlat  <https://orcid.org/0000-0003-0218-0298>

Petra Dieker  <https://orcid.org/0000-0003-3468-4810>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Adamowicz SJ, Boatwright JS, Chain F, Fisher BL, Hogg ID, Leese F, Lijtmaer DA, Mwale M, Naaum AM, Pochon X, Steinke D, Wilson JJ, Wood S, Xu J, Xu S, Zhou X, van der Bank M (2019) Trends in DNA barcoding and metabarcoding. *Genome* 62(3): v–viii. <https://doi.org/10.1139/gen-2019-0054>
- Alberdi A, Gilbert MTP (2019) A guide to the application of Hill numbers to DNA-based diversity analyses. *Molecular Ecology Resources* 19(4): 804–817. <https://doi.org/10.1111/1755-0998.13014>
- Alberdi A, Aizpurua O, Gilbert MTP, Bohmann K, Mahon A (2018) Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution* 9(1): 134–147. <https://doi.org/10.1111/2041-210X.12849>
- Andriollo T, Gillet F, Michaux JR, Ruedi M (2019) The menu varies with metabarcoding practices: A case study with the bat *Plecotus auritus*. *PLoS ONE* 14(7): e0219135. <https://doi.org/10.1371/journal.pone.0219135>
- Andujar C, Arribas P, Yu DW, Vogler AP, Emerson BC (2018) Why the COI barcode should be the community DNA metabarcode for the metazoa. *Molecular Ecology* 27(20): 3968–3975. <https://doi.org/10.1111/mec.14844>
- Andujar C, Creedy TJ, Arribas P, Lopez H, Salces-Castellano A, Perez-Delgado AJ, Vogler AP, Emerson BC (2021) Validated removal of nuclear pseudogenes and sequencing artefacts from mitochondrial metabarcode data. *Molecular Ecology Resources* 21(6): 1772–1787. <https://doi.org/10.1111/1755-0998.13337>
- Ärje J, Melvad C, Jeppesen MR, Madsen SA, Raitoharju J, Rasmussen MS, Iosifidis A, Tirronen V, Gabbouj M, Meissner K, Høye TT, Kotze DJ (2020) Automatic image-based identification and biomass estimation of invertebrates. *Methods in Ecology and Evolution* 11(8): 922–931. <https://doi.org/10.1111/2041-210X.13428>
- Arribas P, Andujar C, Salces-Castellano A, Emerson BC, Vogler AP (2021) The limited spatial scale of dispersal in soil arthropods revealed with whole-community haplotype-level metabarcoding. *Molecular Ecology* 30(1): 48–61. <https://doi.org/10.1111/mec.15591>
- Arstingstall KA, DeBano SJ, Li X, Wooster DE, Rowland MM, Burrows S, Frost K (2023) Investigating the use of pollen DNA metabarcoding to quantify bee foraging and effects of threshold selection. *PLoS ONE* 18(4): e0282715. <https://doi.org/10.1371/journal.pone.0282715>
- Aylagas E, Borja A, Rodriguez-Ezpeleta N (2014) Environmental status assessment using DNA metabarcoding: Towards a genetics based Marine Biotic Index (gAMBI). *PLoS ONE* 9(3): e90529. <https://doi.org/10.1371/journal.pone.0090529>

- Baksay S, Pornon A, Burrus M, Mariette J, Andalo C, Escaravage N (2020) Experimental quantification of pollen with DNA metabarcoding using ITS1 and trnL. *Scientific Reports* 10(1): 4202–4202. <https://doi.org/10.1038/s41598-020-61198-6>
- Barnes MA, Turner CR (2016) The ecology of environmental DNA and implications for conservation genetics. *Conservation Genetics* 17(1): 1–17. <https://doi.org/10.1007/s10592-015-0775-4>
- Batovska J, Piper AM, Valenzuela I, Cunningham JP, Blacket MJ (2021) Developing a non-destructive metabarcoding protocol for detection of pest insects in bulk trap catches. *Scientific Reports* 11(1): 7946. <https://doi.org/10.1038/s41598-021-85855-6>
- Beentjes KK, Speksnijder AGCL, Schilthuizen M, Hoogeveen M, Pastoor R, van der Hoorn BB (2019) Increased performance of DNA metabarcoding of macroinvertebrates by taxonomic sorting. *PLoS ONE* 14(12): e0226527. <https://doi.org/10.1371/journal.pone.0226527>
- Bell KL, de Vere N, Keller A, Richardson RT, Gous A, Burgess KS, Brosi BJ (2016) Pollen DNA barcoding: Current applications and future prospects. *Genome* 59(9): 629–640. <https://doi.org/10.1139/gen-2015-0200>
- Bell KL, Burgess KS, Botsch JC, Dobbs EK, Read TD, Brosi BJ (2019) Quantitative and qualitative assessment of pollen DNA metabarcoding using constructed species mixtures. *Molecular Ecology* 28(2): 431–455. <https://doi.org/10.1111/mec.14840>
- Besson M, Alison J, Bjerger K, Gorochowski TE, Hoyer TT, Jucker T, Mann HMR, Clements CF (2022) Towards the fully automated monitoring of ecological communities. *Ecology Letters* 25(12): 2753–2775. <https://doi.org/10.1111/ele.14123>
- Birk S, Bonne W, Borja A, Brucet S, Courrat A, Poikane S, Solimini A, van de Bund W, Zampoukas N, Hering D (2012) Three hundred ways to assess Europe's surface waters: An almost complete overview of biological methods to implement the Water Framework Directive. *Ecological Indicators* 18: 31–41. <https://doi.org/10.1016/j.ecolind.2011.10.009>
- Blancher P, Lefrançois E, Rimet F, Vasselon V, Argillier C, Arle J, Beja P, Boets P, Boughaba J, Chauvin C, Deacon M, Duncan W, Ejdung G, Erba S, Ferrari B, Fischer H, Hänfling B, Haldin M, Hering D, Hette-Tronquart N, Hiley A, Järvinen M, Jeannot B, Kahlert M, Kelly M, Kleinteich J, Koyuncuoğlu S, Krenek S, Langhein-Winther S, Leese F, Mann D, Marcel R, Marcheggiani S, Meissner K, Mergen P, Monnier O, Narendja F, Neu D, Onofre Pinto V, Pawlowska A, Pawlowski J, Petersen M, Poikane S, Pont D, Renevier M-S, Sandoy S, Svensson J, Trobajo R, Tünde Zagyva A, Tziortzis I, van der Hoorn B, Vasquez MI, Walsh K, Weigand A, Bouchez A (2022) A strategy for successful integration of DNA-based methods in aquatic monitoring. *Metabarcoding and Metagenomics* 6: e85652. <https://doi.org/10.3897/mbmg.6.85652>
- Braukmann TWA, Ivanova NV, Prosser SWJ, Elbrecht V, Steinke D, Ratnasingham S, de Waard JR, Sones JE, Zakharov EV, Hebert PDN (2019) Metabarcoding a diverse arthropod mock community. *Molecular Ecology Resources* 19(3): 711–727. <https://doi.org/10.1111/1755-0998.13008>
- Bruce K, Keskin E (2021) Field and Lab Protocols: Achievements of the DNAqua-Net Working Group 3. *ARPHA Conference Abstracts* 4: e65306. <https://doi.org/10.3897/aca.4.e65306>
- Bruce K, Blackman R, Bourlat SJ, Hellström AM, Bakker J, Bista I, Bohmann K, Bouchez A, Brys R, Clark K, Elbrecht V, Fazi S, Fonseca V, Hänfling B, Leese F, Mächler E, Mahon AR, Meissner K, Panksep K, Pawlowski J, Schmidt Yáñez P, Seymour M, Thälinger B,

- Valentini A, Woodcock P, Traugott M, Vasselon V, Deiner K (2021) A practical guide to DNA-based methods for biodiversity assessment. Advanced Books. <https://doi.org/10.3897/ab.e68634>
- Buchner D, Beermann AJ, Laini A, Rolauffs P, Vitecek S, Hering D, Leese F (2019) Analysis of 13,312 benthic invertebrate samples from German streams reveals minor deviations in ecological status class between abundance and presence/absence data. PLoS ONE 14(12): e0226547. <https://doi.org/10.1371/journal.pone.0226547>
- Buchner D, Macher TH, Beermann AJ, Werner MT, Leese F (2021) Standardized high-throughput biomonitoring using DNA metabarcoding: Strategies for the adoption of automated liquid handlers. Environmental Science and Ecotechnology 8: 100122. <https://doi.org/10.1016/j.esec.2021.100122>
- Buchner D, Beermann A, Hörrn T, Enss J, Frenzel M, Li Y, Müller J, Pauls SU, Sorg M, Haase P, Leese F (2023) German-wide Malaise trap metabarcoding estimates over 33,000 insect species. BioRxiv. <https://doi.org/10.1101/2023.05.04.539402>
- Carew ME, Coleman RA, Hoffmann AA (2018) Can non-destructive DNA extraction of bulk invertebrate samples be used for metabarcoding? PeerJ 6: e4980. <https://doi.org/10.7717/peerj.4980>
- Castalanelli MA, Severtson DL, Brumley CJ, Szito A, Footitt RG, Grimm M, Munyard KG, David M (2010) A rapid non-destructive DNA extraction method for insects and other arthropods. Journal of Asia-Pacific Entomology 13(3): 243–248. <https://doi.org/10.1016/j.aspen.2010.04.003>
- CBOL Plant Working Group, Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, van der Bank M, Chase MW, Cowan RS, Erickson DL, Fazekas AJ, Graham SW, James KE, Kim K-J, Kress WJ, Schneider H, van AlphenStahl J, Barrett SCH, van den Berg C, Bogarin D, Burgess KS, Cameron KM, Carine M, Chacón J, Clark A, Clarkson JJ, Conrad F, Devey DS, Ford CS, Hedderson TAJ, Hollingsworth ML, Husband BC, Kelly LJ, Kesanakurti PR, Kim JS, Kim Y-D, Lahaye R, Lee H-L, Long DG, Madriñán S, Maurin O, Meusnier I, Newmaster SG, Park C-W, Percy DM, Petersen G, Richardson JE, Salazar GA, Savolainen V, Seberg O, Wilkinson MJ, Yi D-K, Little DP (2009) A DNA barcode for land plants. Proceedings of the National Academy of Sciences of the United States of America 106(31): 12794–12797. <https://doi.org/10.1073/pnas.0905845106>
- Clarke LJ, Soubrier J, Weyrich LS, Cooper A (2014) Environmental metabarcodes for insects: In silico PCR reveals potential for taxonomic bias. Molecular Ecology Resources 14(6): 1160–1170. <https://doi.org/10.1111/1755-0998.12265>
- Compson ZG, McClenaghan B, Singer GAC, Fahner NA, Hajibabaei M (2020) Metabarcoding From Microbes to Mammals: Comprehensive Bioassessment on a Global Scale. Frontiers in Ecology and Evolution 8: 581835. <https://doi.org/10.3389/fevo.2020.581835>
- Cordier T, Alonso-Saez L, Apotheloz-Perret-Gentil L, Aylagas E, Bohan DA, Bouchez A, Chariton A, Creer S, Fruhe L, Keck F, Keeley N, Laroche O, Leese F, Pochon X, Stoeck T, Pawlowski J, Lanzen A (2021) Ecosystems monitoring powered by environmental genomics: A review of current strategies with an implementation roadmap. Molecular Ecology 30(13): 2937–2958. <https://doi.org/10.1111/mec.15472>
- Creedy T, Andujar C, Meramveliotakis E, Noguerales V, Overcast I, Papadopoulou A, Morlon H, Vogler A, Emerson B, Arribas P (2021) Coming of age for COI metabarcoding of whole organism community DNA: towards bioinformatic harmonisation. Authorea Preprints: under review. <https://doi.org/10.22541/au.162141276.61766048/v1>
- Cuff JP, Windsor FM, Tercel MPTG, Kitson JJN, Evans DM (2022) Overcoming the pitfalls of merging dietary metabarcoding into ecological networks. Methods in Ecology and Evolution 13(3): 545–559. <https://doi.org/10.1111/2041-210X.13796>

- da Silva LP, Mata VA, Lopes PB, Pereira P, Jarman SN, Lopes RJ, Beja P (2019) Advancing the integration of multi-marker metabarcoding data in dietary analysis of trophic generalists. *Molecular Ecology Resources* 19(6): 1420–1432. <https://doi.org/10.1111/1755-0998.13060>
- Darby BJ, Todd TC, Herman MA (2013) High-throughput amplicon sequencing of rRNA genes requires a copy number correction to accurately reflect the effects of management practices on soil nematode community structure. *Molecular Ecology* 22(21): 5456–5471. <https://doi.org/10.1111/mec.12480>
- Darby B, Bryant R, Keller A, Jochim M, Moe J, Schreiner Z, Pratt C, Euliss NH, Park M, Simmons R, Otto C (2020) Molecular sequencing and morphological identification reveal similar patterns in native bee communities across public and private grasslands of eastern North Dakota. *PLoS ONE* 15(1): e0227918. <https://doi.org/10.1371/journal.pone.0227918>
- de Kerdrel GA, Andersen JC, Kennedy SR, Gillespie R, Krehenwinkel H (2020) Rapid and cost-effective generation of single specimen multilocus barcoding data from whole arthropod communities by multiple levels of multiplexing. *Scientific Reports* 10(1): 78. <https://doi.org/10.1038/s41598-019-54927-z>
- Deagle BE, Thomas AC, McInnes JC, Clarke LJ, Vesterinen EJ, Clare EL, Kartzinel TR, Eveson JP (2019) Counting with DNA in metabarcoding studies: How should we convert sequence reads to dietary data? *Molecular Ecology* 28(2): 391–406. <https://doi.org/10.1111/mec.14734>
- Díaz S, Settele J, Brondízio E, Ngo HT, Guèze M, Agard J, Arneth A, Balvanera P, Brauman K, Butchart S, Chan K, Garibaldi L, Ichii K, Liu J, Subramanian SM, Midgley G, Miloslavich P, Molnár Z, Obura D, Pfaff A, Polasky S, Purvis A, Razzaque J, Reyers B, Chowdhury RR, Shin Y-J, Visseren-Hamakers I, Willis K, Zayas C (2020) Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
- Dickie IA, Boyer S, Buckley HL, Duncan RP, Gardner PP, Hogg ID, Holdaway RJ, Lear G, Makiola A, Morales SE, Powell JR, Weaver L (2018) Towards robust and repeatable sampling methods in eDNA-based studies. *Molecular Ecology Resources* 18(5): 940–952. <https://doi.org/10.1111/1755-0998.12907>
- Directive 2000/60/EC (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 Establishing a framework for Community action in the field of water policy. *Official Journal of the European Union L 327*: 1–73.
- Directive 2008/56/EC (2008) Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for Community action in the field of marine environmental policy. *Official Journal of the European Union L 164*: 19–40.
- Doi H, Fukaya K, Oka SI, Sato K, Kondoh M, Miya M (2019) Evaluation of detection probabilities at the water-filtering and initial PCR steps in environmental DNA metabarcoding using a multispecies site occupancy model. *Scientific Reports* 9(1): 3581. <https://doi.org/10.1038/s41598-019-40233-1>
- Dunker S, Motivans E, Rakosy D, Boho D, Mäder P, Hornick T, Knight TM (2020) Pollen analysis using multispectral imaging flow cytometry and deep learning. *The New Phytologist* 229(1): 593–606. <https://doi.org/10.1111/nph.16882>
- Elbrecht V, Leese F (2015) Can DNA-based ecosystem assessments quantify species abundance? Testing primer bias and biomass-sequence relationships with an innovative metabarcoding protocol. *PLoS ONE* 10(7): e0130324. <https://doi.org/10.1371/journal.pone.0130324>

- Elbrecht V, Steinke D (2019) Scaling up DNA metabarcoding for freshwater macrozoobenthos monitoring. *Freshwater Biology* 64: 380–387. <https://doi.org/10.1111/fwb.13220>
- Elbrecht V, Peinert B, Leese F (2017) Sorting things out: Assessing effects of unequal specimen biomass on DNA metabarcoding. *Ecology and Evolution* 7(17): 6918–6926. <https://doi.org/10.1002/ece3.3192>
- Elbrecht V, Vamos EE, Steinke D, Leese F (2018) Estimating intraspecific genetic diversity from community DNA metabarcoding data. *PeerJ* 6: e4644. <https://doi.org/10.7717/peerj.4644>
- Elbrecht V, Braukmann TWA, Ivanova NV, Prosser SWJ, Hajibabaei M, Wright M, Zakharov EV, Hebert PDN, Steinke D (2019) Validation of COI metabarcoding primers for terrestrial arthropods. *PeerJ* 7: e7745. <https://doi.org/10.7717/peerj.7745>
- Elbrecht V, Bourlat SJ, Hörren T, Lindner A, Mordente A, Noll NW, Sorg M, Zizka VMA (2020) Pooling size sorted malaise trap fractions to maximise taxon recovery with metabarcoding. *BioRxiv*. <https://doi.org/10.1101/2020.06.09.118950>
- Esnaola A, Arrizabalaga-Escudero A, González-Esteban J, Elosegi A, Aihartza J (2018) Determining diet from faeces: Selection of metabarcoding primers for the insectivore Pyrenean desman (*Galemys pyrenaicus*). *PLoS ONE* 13(12): e0208986. <https://doi.org/10.1371/journal.pone.0208986>
- Fernandes K, van der Heyde M, Coghlan M, Wardell-Johnson G, Bunce M, Harris R, Nevill P (2019) Invertebrate DNA metabarcoding reveals changes in communities across mine site restoration chronosequences. *Restoration Ecology* 27(5): 1177–1186. <https://doi.org/10.1111/rec.12976>
- Fields B, Moeskjær S, Friman V-P, Andersen SU, Young JPW (2020) MAUI-seq: Metabarcoding using amplicons with unique molecular identifiers to improve error correction. *BioRxiv*. <https://doi.org/10.21203/rs.2.21630/v1>
- Formenti G, Theissinger K, Fernandes C, Bista I, Bombarely A, Bleidorn C, Ciofi C, Crottini A, Godoy JA, Høglund J, Malukiewicz J, Mouton A, Oomen RA, Paez S, Palsboll PJ, Pampoulie C, Ruiz-Lopez MJ, Svoldal H, Theofanopoulou C, de Vries J, Waldvogel AM, Zhang G, Mazzoni CJ, Jarvis ED, Balint M, Formenti G, Theissinger K, Fernandes C, Bista I, Bombarely A, Bleidorn C, Čiampor F, Ciofi C, Crottini A, Godoy JA, Høglund J, Malukiewicz J, Mouton A, Oomen RA, Paez S, Palsbøll P, Pampoulie C, Ruiz-López MJ, Svoldal H, Theofanopoulou C, de Vries J, Waldvogel A-M, Zhang G, Mazzoni CJ, Jarvis E, Bálint M, Aghayan SA, Alioto TS, Almudi I, Alvarez N, Alves PC, Amorim IR, Antunes A, Arribas P, Baldrian P, Berg PR, Bertorelle G, Böhne A, Bonisoli-Alquati A, Boštjančić LL, Boussau B, Breton CM, Buzan E, Campos PF, Carreras C, Castro LFI, Chueca LJ, Conti E, Cook-Deegan R, Croll D, Cunha MV, Delsuc F, Dennis AB, Dimitrov D, Faria R, Favre A, Fedrigo OD, Fernández R, Ficetola GF, Flot J-F, Gabaldón T, Galea Agius DR, Gallo GR, Giani AM, Gilbert MTP, Grebenc T, Guschanski K, Guyot R, Hausdorf B, Hawlitschek O, Heintzman PD, Heinze B, Hiller M, Husemann M, Iannucci A, Irisarri I, Jakobsen KS, Jentoft S, Klinga P, Kloch A, Kratochwil CF, Kusche H, Layton KKS, Leonard JA, Lerat E, Liti G, Manousaki T, Marques-Bonet T, Matos-Maraví P, Matschiner M, Maumus F, Mc Cartney AM, Meiri S, Melo-Ferreira J, Mengual X, Monaghan MT, Montagna M, Mysłajek RW, Neiber MT, Nicolas V, Novo M, Ozretić P, Palero F, Pârvulescu L, Pascual M, Paulo OS, Pavlek M, Pegueroles C, Pellissier L, Pesole G, Primmer CR, Riesgo A, Rüber L, Rubolini D, Salvi D, Seehausen O, Seidel M, Secomandi S, Studer B, Theodoridis S, Thines M, Urban L, Vasemägi A, Vella A, Vella N, Vernes SC, Vernesi C, Vieites DR, Waterhouse RM, Wheat CW, Wörheide G, Wurm Y, Zammit G (2022) The era of reference genomes in conservation genomics. *Trends in Ecology & Evolution* 37(3): 197–202. <https://doi.org/10.1016/j.tree.2021.11.008>

- Frantz A, Schaul M, Pope LC, Fack F, Schley L, Muller CP, Roper TM (2004) Estimating population size by genotyping remotely plucked hair: The Eurasian badger. *Journal of Applied Ecology* 41(5): 985–995. <https://doi.org/10.1111/j.0021-8901.2004.00951.x>
- Frøslev TG, Kjølner R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, Hansen AJ (2017) Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nature Communications* 8(1): 1188. <https://doi.org/10.1038/s41467-017-01312-x>
- Fukaya K, Murakami H, Yoon S, Minami K, Osada Y, Yamamoto S, Masuda R, Kasai A, Miyashita K, Minamoto T, Kondoh M (2020) Estimating fish population abundance by integrating quantitative data on environmental DNA and hydrodynamic modelling. *Molecular Ecology* 30(13): 3057–3067. <https://doi.org/10.1111/mec.15530>
- Garrido-Sanz L, Senar MÀ, Piñol J (2020) Estimation of the relative abundance of species in artificial mixtures of insects using low-coverage shotgun metagenomics. *Metabarcoding and Metagenomics* 4: e48281. <https://doi.org/10.3897/mbmg.4.48281>
- Giebner H, Langen K, Bourlat SJ, Kukowka S, Mayer C, Astrin JJ, Misof B, Fonseca VG (2020) Comparing diversity levels in environmental samples: DNA sequence capture and metabarcoding approaches using 18S and COI genes. *Molecular Ecology Resources* 20(5): 1333–1345. <https://doi.org/10.1111/1755-0998.13201>
- Gil V, Pinho CJ, Aguiar CAS, Jardim C, Rebelo R, Vasconcelos R (2020) Questioning the proverb ‘more haste, less speed’: Classic versus metabarcoding approaches for the diet study of a remote island endemic gecko. *PeerJ* 8: e8084. <https://doi.org/10.7717/peerj.8084>
- Goberna M, Verdu M (2016) Predicting microbial traits with phylogenies. *The ISME Journal* 10(4): 959–967. <https://doi.org/10.1038/ismej.2015.171>
- Gueuning M, Ganser D, Blaser S, Albrecht M, Knop E, Praz C, Frey JE (2019) Evaluating next-generation sequencing (NGS) methods for routine monitoring of wild bees: Metabarcoding, mitogenomics or NGS barcoding. *Molecular Ecology Resources* 19(4): 847–862. <https://doi.org/10.1111/1755-0998.13013>
- Haase P, Lohse SPS, Schindehütte KSA, Rolauffs P, Hering D (2004) Assessing streams in Germany with benthic invertebrates: Development of a practical standardised protocol for macroinvertebrate sampling and sorting. *Limnologia* 34(4): 349–365. [https://doi.org/10.1016/S0075-9511\(04\)80005-7](https://doi.org/10.1016/S0075-9511(04)80005-7)
- Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W, Müller A, Sumser H, Hörrn T, Goulson D, de Kroon H (2017) More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS ONE* 12(10): e0185809. <https://doi.org/10.1371/journal.pone.0185809>
- Harrison JG, Randolph GD, Buerkle CA (2021) Characterizing microbiomes via sequencing of marker loci: Techniques to improve throughput, account for crosscontamination, and reduce cost. *mSystems* 6(4): e00294–e00221. <https://doi.org/10.1128/mSystems.00294-21>
- Hausmann A, Segerer AH, Greifstein T, Knubben J, Morinière J, Bozicevic V, Doczkal D, Günter A, Ulrich W, Habel JC (2020) Toward a standardized quantitative and qualitative insect monitoring scheme. *Ecology and Evolution* 10(9): 4009–4020. <https://doi.org/10.1002/ece3.6166>
- Hawkins J, de Vere N, Griffith A, Ford CR, Allainguillaume J, Hegarty MJ, Baillie L, Adams-Groom B (2015) Using DNA Metabarcoding to Identify the Floral Composition of Honey: A New Tool for Investigating Honey Bee Foraging Preferences. *PLoS ONE* 10(8): e0134735. <https://doi.org/10.1371/journal.pone.0134735>
- Hope PR, Bohmann K, Gilbert MTP, Zepeda-Mendoza ML, Razgour O, Jones G (2014) Second generation sequencing and morphological faecal analysis reveal unexpected

- foraging behaviour by *Myotis nattereri* (Chiroptera, Vespertilionidae) in winter. *Frontiers in Zoology* 11(1): 39. <https://doi.org/10.1186/1742-9994-11-39>
- Hoshino T, Inagaki F (2017) Application of Stochastic Labeling with Random-Sequence Barcodes for Simultaneous Quantification and Sequencing of Environmental 16S rRNA Genes. *PLoS ONE* 12(1): e0169431. <https://doi.org/10.1371/journal.pone.0169431>
- Hoshino T, Nakao R, Doi H, Minamoto T (2021) Simultaneous absolute quantification and sequencing of fish environmental DNA in a mesocosm by quantitative sequencing technique. *Scientific Reports* 11(1): 4372. <https://doi.org/10.1038/s41598-021-83318-6>
- Høye TT, Årje J, Bjerger K, Hansen OLP, Iosifidis A, Leese F, Mann HMR, Meissner K, Melvad C, Raitoharju J (2021) Deep learning and computer vision will transform entomology. *Proceedings of the National Academy of Sciences of the United States of America* 118(2): e2002545117. <https://doi.org/10.1073/pnas.2002545117>
- Iwaszkiewicz-Eggebrecht E, Granqvist E, Buczek M, Prus M, Roslin T, Tack AJM, Andersson AF, Miraldo A, Ronquist F, Łukasik P (2022) Optimizing insect metabarcoding using replicated mock communities. *BioRxiv*. <https://doi.org/10.1101/2022.06.20.496906>
- Jabara CB, Jones CD, Roach J, Anderson JA, Swanstrom R (2011) Accurate sampling and deep sequencing of the HIV-1 protease gene using a Primer ID. *Proceedings of the National Academy of Sciences of the United States of America* 108(50): 20166–20171. <https://doi.org/10.1073/pnas.1110064108>
- Ji Y, Huotari T, Roslin T, Schmidt NM, Wang J, Yu DW, Ovaskainen O (2020) SPIKEPIPE: A metagenomic pipeline for the accurate quantification of eukaryotic species occurrences and intraspecific abundance change using DNA barcodes or mitogenomes. *Molecular Ecology Resources* 20(1): 256–267. <https://doi.org/10.1111/1755-0998.13057>
- Jo T, Takao K, Minamoto T (2022) Linking the state of environmental DNA to its application for biomonitoring and stock assessment: Targeting mitochondrial/nuclear genes, and different DNA fragment lengths and particle sizes. *Environmental DNA* 4(2): 271–283. <https://doi.org/10.1002/edn3.253>
- Jusino MA, Banik MT, Palmer JM, Wray AK, Xiao L, Pelton E, Barber JR, Kawahara AY, Gratton C, Peery MZ, Lindner DL (2018) An improved method for utilizing high-throughput amplicon sequencing to determine the diets of insectivorous animals. *Molecular Ecology Resources* 19(1): 176–190. <https://doi.org/10.1111/1755-0998.12951>
- Keller A, Danner N, Grimmer G, Ankenbrand M, von der Ohe K, von der Ohe W, Rost S, Härtel S, Steffan-Dewenter I (2015) Evaluating multiplexed next-generation sequencing as a method in palynology for mixed pollen samples. *Plant Biology* 17(2): 558–566. <https://doi.org/10.1111/plb.12251>
- Kelly RP, Shelton AO, Gallego R (2019) Understanding PCR processes to draw meaningful conclusions from environmental DNA studies. *Scientific Reports* 9(1): 12133. <https://doi.org/10.1038/s41598-019-48546-x>
- Kelly RP, Lodge DM, Lee KN, Theroux S, Sepulveda AJ, Scholin CA, Craine JM, Andruszkiewicz Allan E, Nichols KM, Parsons KM, Goodwin KD, Gold Z, Chavez FP, Noble RT, Abbott CL, Baerwald MR, Naaum AM, Thielen PM, Simons AL, Jerde CL, Duda JJ, Hunter ME, Hagan JA, Meyer RS, Steele JA, Stoeckle MY, Bik HM, Meyer CP, Stein E, James KE, Thomas AC, Demir-Hilton E, Timmers MA, Griffith JF, Weise MJ, Weisberg SB (2023) Toward a national eDNA strategy for the United States. *Environmental DNA* 00: 1–10. <https://doi.org/10.1002/edn3.432>
- Kinde I, Wu J, Papadopoulos N, Kinzler KW, Vogelstein B (2011) Detection and quantification of rare mutations with massively parallel sequencing. *Proceedings of the National Academy of Sciences of the United States of America* 108(23): 9530–9535. <https://doi.org/10.1073/pnas.1105422108>

- Kirse A, Bourlat SJ, Langen K, Fonseca VG (2021) Unearthing the Potential of Soil eDNA Metabarcoding – Towards Best Practice Advice for Invertebrate Biodiversity Assessment. *Frontiers in Ecology and Evolution* 9: 630560. <https://doi.org/10.3389/fevo.2021.630560>
- Kirse A, Bourlat SJ, Langen K, Zapke B, Zizka VMA (2023) Comparison of destructive and nondestructive DNA extraction methods for the metabarcoding of arthropod bulk samples. *Molecular Ecology Resources* 23(1): 92–105. <https://doi.org/10.1111/1755-0998.13694>
- Kivioja T, Vähärautio A, Karlsson K, Bonke M, Enge M, Linnarsson S, Taipale J (2012) Counting absolute numbers of molecules using unique molecular identifiers. *Nature Methods* 9(1): 72–74. <https://doi.org/10.1038/nmeth.1778>
- Kolter AG, Gemeinholzer B (2021) Plant DNA barcoding necessitates marker-specific efforts to establish more comprehensive reference databases. *Genome* 64(3): 265–298. <https://doi.org/10.1139/gen-2019-0198>
- Kraaijeveld K, de Weger LA, Ventayol Garcia M, Buermans H, Frank J, Hiemstra PS, den Dunnen JT (2015) Efficient and sensitive identification and quantification of airborne pollen using next-generation DNA sequencing. *Molecular Ecology Resources* 15(1): 8–16. <https://doi.org/10.1111/1755-0998.12288>
- Kratschmer S, Petrović B, Curto M, Meimberg H, Pachinger B (2020) Pollen availability for the Horned mason bee (*Osmia cornuta*) in regions of different land use and landscape structures. *Ecological Entomology* 45(3): 525–537. <https://doi.org/10.1111/een.12823>
- Krehenwinkel H, Kennedy S, Pekár S, Gillespie RG (2017a) A cost-efficient and simple protocol to enrich prey DNA from extractions of predatory arthropods for large-scale gut content analysis by Illumina sequencing. *Methods in Ecology and Evolution* 8(1): 126–134. <https://doi.org/10.1111/2041-210X.12647>
- Krehenwinkel H, Wolf M, Lim JY, Rominger AJ, Simison WB, Gillespie RG (2017b) Estimating and mitigating amplification bias in qualitative and quantitative arthropod metabarcoding. *Scientific Reports* 7(1): 17668. <https://doi.org/10.1038/s41598-017-17333-x>
- Krehenwinkel H, Fong M, Kennedy S, Huang EG, Noriyuki S, Cayetano L, Gillespie R (2018) The effect of DNA degradation bias in passive sampling devices on metabarcoding studies of arthropod communities and their associated microbiota. *PLoS ONE* 13(1): e0189188. <https://doi.org/10.1371/journal.pone.0189188>
- Krehenwinkel H, Pomerantz A, Henderson JB, Kennedy SR, Lim JY, Swamy V, Shoobridge JD, Graham N, Patel NH, Gillespie RG, Prost S (2019) Nanopore sequencing of long ribosomal DNA amplicons enables portable and simple biodiversity assessments with high phylogenetic resolution across broad taxonomic scale. *GigaScience* 8(5): 1–16. <https://doi.org/10.1093/gigascience/giz006>
- Lafage D, Elbrecht V, Cuff JP, Steinke D, Hambäck PA, Erlandsson A (2019) A new primer for metabarcoding of spider gut contents. *Environmental DNA* 2(2): 234–243. <https://doi.org/10.1002/edn3.62>
- Lamb PD, Hunter E, Pinnegar JK, Creer S, Davies RG, Taylor MI (2019) How quantitative is metabarcoding: A meta-analytical approach. *Molecular Ecology* 28(2): 420–430. <https://doi.org/10.1111/mec.14920>
- Leese F, Altermatt F, Bouchez A, Ekrem T, Hering D, Meissner K, Mergen P, Pawlowski J, Pigott J, Rimet F, Steinke D, Taberlet P, Weigand A, Abarenkov K, Beja P, Bervoets L, Björnsdóttir S, Boets P, Boggero A, Bones A, Borja Á, Bruce K, Bursić V, Carlsson J, Čiampor F, Čiamporová-Zatovičová Z, Coissac E, Costa F, Costache M, Creer S, Csabai Z, Deiner K, DelValls Á, Drakare S, Duarte S, Eleršek T, Fazi S, Fišer C, Flot J-F, Fonseca V, Fontaneto D, Grabowski M, Graf W, Guðbrandsson J, Hellström M, Hershkovitz Y, Hollingsworth

- P, Japoshvili B, Jones J, Kahlert M, Kalamujic Stroil B, Kasapidis P, Kelly M, Kelly-Quinn M, Keskin E, Kõljalg U, Ljubešić Z, Maček I, Mächler E, Mahon A, Marečková M, Mejdandzic M, Mircheva G, Montagna M, Moritz C, Mulk V, Naumoski A, Navodaru I, Padisák J, Pálsson S, Panksep K, Penev L, Petrusek A, Pfannkuchen M, Primmer C, Rinkevich B, Rotter A, Schmidt-Kloiber A, Segurado P, Speksnijder A, Stoev P, Strand M, Šulčius S, Sundberg P, Traugott M, Tsigenopoulos C, Turon X, Valentini A, van der Hoorn B, Várбірó G, Vasquez Hadjilyra M, Viguri J, Vitonytė I, Vogler A, Vrålstad T, Wägele W, Wenne R, Winding A, Woodward G, Zegura B, Zimmermann J (2016) DNAqua-Net: Developing new genetic tools for bioassessment and monitoring of aquatic ecosystems in Europe. *Research Ideas and Outcomes* 2: e11321. <https://doi.org/10.3897/rio.2.e11321>
- Leese F, Bouchez A, Abarenkov K, Altermatt F, Borja Á, Bruce K, Ekrem T, Čiampor Jr F, Čiamporová-Zaťovičová Z, Costa FO, Duarte S, Elbrecht V, Fontaneto D, Franc A, Geiger MF, Hering D, Kahlert M, Kalamujic Stroil B, Kelly M, Keskin E, Liska I, Mergen P, Meissner K, Pawlowski J, Penev L, Reyjol Y, Rotter A, Steinke D, van der Wal B, Vitecek S, Zimmermann J, Weigand AM (2018) Why We Need Sustainable Networks Bridging Countries, Disciplines, Cultures and Generations for Aquatic Biomonitoring 2.0: A Perspective Derived From the DNAqua-Net COST Action. *Advances in Ecological Research* 58(Part 1): 63–99. <https://doi.org/10.1016/bs.aecr.2018.01.001>
- Leese F, Woppowa L, Bálint M, Höss S, Krehenwinkel H, Lötters S, Meissner K, Nowak C, Rausch P, Rduch V, Rulik B, Weigand AM, Zimmermann J, Koschorreck J, Züghart W (2023) DNA-basierte Biodiversitätsanalysen im Natur- und Umweltschutz: Welche Optionen haben wir für eine Standardisierung? – Eine Handlungsempfehlung aus Forschung und Praxis. *BfN-Schriften* 666.
- Leidenfrost RM, Bänsch S, Prudnikow L, Brenig B, Westphal C, Wünschiers R (2020) Analyzing the dietary diary of Bumble Bee. *Frontiers in Plant Science* 11: 287. <https://doi.org/10.3389/fpls.2020.00287>
- Leray M, Knowlton N (2017) Random sampling causes the low reproducibility of rare eukaryotic OTUs in Illumina COI metabarcoding. *PeerJ* 5: e3006. <https://doi.org/10.7717/peerj.3006>
- Lewin HA, Richards S, Lieberman Aiden E, Allende ML, Archibald JM, Balint M, Barker KB, Baumgartner B, Belov K, Bertorelle G, Blaxter ML, Cai J, Caperello ND, Carlson K, Castilla-Rubio JC, Chaw SM, Chen L, Childers AK, Coddington JA, Conde DA, Corominas M, Crandall KA, Crawford AJ, DiPalma F, Durbin R, Ebenezer TE, Edwards SV, Fedrigo O, Flicek P, Formenti G, Gibbs RA, Gilbert MTP, Goldstein MM, Graves JM, Greely HT, Grigoriev IV, Hackett KJ, Hall N, Haussler D, Helgen KM, Hogg CJ, Isobe S, Jakobsen KS, Janke A, Jarvis ED, Johnson WE, Jones SJM, Karlsson EK, Kersey PJ, Kim JH, Kress WJ, Kuraku S, Lawniczak MKN, Leebens-Mack JH, Li X, Lindblad-Toh K, Liu X, Lopez JV, Marques-Bonet T, Mazar S, Mazet JAK, Mazzoni CJ, Myers EW, O'Neill RJ, Paez S, Park H, Robinson GE, Roquet C, Ryder OA, Sabir JSM, Shaffer HB, Shank TM, Sherkow JS, Soltis PS, Tang B, Tedersoo L, Uliano-Silva M, Wang K, Wei X, Wetzler R, Wilson JL, Xu X, Yang H, Yoder AD, Zhang G (2022) The Earth BioGenome Project 2020: Starting the clock. *Proceedings of the National Academy of Sciences of the United States of America* 119(4): e2115635118. <https://doi.org/10.1073/pnas.2115635118>
- Lister BC, Garcia A (2018) Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proceedings of the National Academy of Sciences of the United States of America* 115(44): E10397–E10406. <https://doi.org/10.1073/pnas.1722477115>
- Liu M, Clarke LJ, Baker SC, Jordan GJ, Burridge CP (2020) A practical guide to DNA metabarcoding for entomological ecologists. *Ecological Entomology* 45(3): 373–385. <https://doi.org/10.1111/een.12831>

- Liu M, Burridge CP, Clarke LJ, Baker SC, Jordan GJ (2023) Does phylogeny explain bias in quantitative DNA metabarcoding? *Metabarcoding and Metagenomics* 7: e101266. <https://doi.org/10.3897/mbmg.7.101266>
- Luo M, Ji Y, Warton D, Yu DW (2023) Extracting abundance information from DNA-based data. *Molecular Ecology Resources* 23(1): 174–189. <https://doi.org/10.1111/1755-0998.13703>
- Macías-Hernández N, Athey K, Tonzo V, Wangenstein OS, Arnedo M, Harwood JD (2018) Molecular gut content analysis of different spider body parts. *PLoS ONE* 13(5): e0196589. <https://doi.org/10.1371/journal.pone.0196589>
- Marquina D, Esparza-Salas R, Roslin T, Ronquist F (2019) Establishing arthropod community composition using metabarcoding: Surprising inconsistencies between soil samples and preservative ethanol and homogenate from Malaise trap catches. *Molecular Ecology Resources* 19(6): 1516–1530. <https://doi.org/10.1111/1755-0998.13071>
- Martoni F, Piper AM, Rodoni BC, Blacket MJ (2022) Disentangling bias for non-destructive insect metabarcoding. *PeerJ* 10: e12981. <https://doi.org/10.7717/peerj.12981>
- Matos-Maraví P, Ritter CD, Barnes CJ, Nielsen M, Olsson U, Wahlberg N, Marquina D, Sääksjärvi I, Antonelli A (2019) Biodiversity seen through the perspective of insects: 10 simple rules on methodological choices and experimental design for genomic studies. *PeerJ* 7: e6727. <https://doi.org/10.7717/peerj.6727>
- Meng G, Li Y, Yang C, Liu S (2019) MitoZ: A toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic Acids Research* 47(11): e63. <https://doi.org/10.1093/nar/gkz173>
- Meusnier I, Singer GA, Landry JF, Hickey DA, Hebert PD, Hajibabaei M (2008) A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics* 9(1): 214. <https://doi.org/10.1186/1471-2164-9-214>
- Milla L, Sniderman K, Lines R, Mousavi-Derazmahalleh M, Encinas-Viso F (2021) Pollen DNA metabarcoding identifies regional provenance and high plant diversity in Australian honey. *Ecology and Evolution* 11(13): 8683–8698. <https://doi.org/10.1002/ece3.7679>
- Miotto RA, Rodrigues FP, Ciocheti G, Galetti Jr PM (2007) Determination of the Minimum Population Size of Pumas (*Puma concolor*) Through Fecal DNA Analysis in Two Protected Cerrado Areas in the Brazilian Southeast. *Biotropica* 39(5): 647–654. <https://doi.org/10.1111/j.1744-7429.2007.00315.x>
- Nichols RV, Vollmers C, Newsom LA, Wang Y, Heintzman PD, Leighton M, Green RE, Shapiro B (2018) Minimizing polymerase biases in metabarcoding. *Molecular Ecology Resources* 18(5): 927–939. <https://doi.org/10.1111/1755-0998.12895>
- Palmer JM, Jusino MA, Banik MT, Lindner DL (2018) Non-biological synthetic spike-in controls and the AMPtk software pipeline improve mycobiome data. *PeerJ* 6: e4925. <https://doi.org/10.7717/peerj.4925>
- Paula DP, Timbo RV, Togawa RC, Vogler AP, Andow DA (2023) Quantitative prey species detection in predator guts across multiple trophic levels by mapping unassembled shotgun reads. *Molecular Ecology Resources* 23(1): 64–80. <https://doi.org/10.1111/1755-0998.13690>
- Pawlowski J, Kelly-Quinn M, Altermatt F, Apotheloz-Perret-Gentil L, Beja P, Boggero A, Borja A, Bouchez A, Cordier T, Domaizon I, Feio MJ, Filipe AF, Fornaroli R, Graf W, Herder J, van der Hoorn B, Iwan Jones J, Sagova-Mareckova M, Moritz C, Barquin J, Piggott JJ, Pinna M, Rimet F, Rinkevich B, Sousa-Santos C, Specchia V, Trobajo R, Vasselon V, Vitecek S, Zimmerman J, Weigand A, Leese F, Kahlert M (2018) The future of biotic indices in the ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems. *The Science of the Total Environment* 637–638: 1295–1310. <https://doi.org/10.1016/j.scitotenv.2018.05.002>

- Pedro PM, Amorim J, Rojas MVR, Sa IL, Galardo AKR, Santos Neto NF, Pires de Carvalho D, Nabas Ribeiro KA, Razzolini MTP, Sallum MAM (2020) Culicidae-centric metabarcoding through targeted use of D2 ribosomal DNA primers. *PeerJ* 8: e9057. <https://doi.org/10.7717/peerj.9057>
- Piñol J, Mir G, Gomez-Polo P, Agustí N (2015) Universal and blocking primer mismatches limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular Ecology Resources* 15(4): 819–830. <https://doi.org/10.1111/1755-0998.12355>
- Piñol J, Senar MA, Symondson WOC (2019) The choice of universal primers and the characteristics of the species mixture determine when DNA metabarcoding can be quantitative. *Molecular Ecology* 28(2): 407–419. <https://doi.org/10.1111/mec.14776>
- Piper AM, Batovska J, Cogan NOI, Weiss J, Cunningham JP, Rodoni BC, Blacket MJ (2019) Prospects and challenges of implementing DNA metabarcoding for high-throughput insect surveillance. *GigaScience* 8: 1–22. <https://doi.org/10.1093/gigascience/giz092>
- Pornon A, Escaravage N, Burrus M, Holota H, Khimoun A, Mariette J, Pellizzari C, Iribar A, Etienne R, Taberlet P, Vidal M, Winterton P, Zinger L, Andalo C (2016) Using metabarcoding to reveal and quantify plant-pollinator interactions. *Scientific Reports* 6(1): 27282. <https://doi.org/10.1038/srep27282>
- Pornon A, Andalo C, Burrus M, Escaravage N (2017) DNA metabarcoding data unveils invisible pollination networks. *Scientific Reports* 7(1): 16828. <https://doi.org/10.1038/s41598-017-16785-5>
- Porter TM, Hajibabaei M (2018) Scaling up: A guide to high-throughput genomic approaches for biodiversity analysis. *Molecular Ecology* 27(2): 313–338. <https://doi.org/10.1111/mec.14478>
- Potts SG, Dauber J, Hochkirch A, Oteman B, Roy DB, Ahmé K, Biesmeijer K, Breeze TD, Carvell C, Ferreira C, FitzPatrick Ú, Isaac NJB, Kuussaari M, Ljubomirov T, Maes J, Ngo H, Pardo A, Polce C, Quaranta M, Settele J, Sorg M, Stefanescu C, Vujić A (2021) Proposal for EU Pollinator Monitoring Scheme. JRC Technical Report. Publications Office of the European Union, Luxembourg, 313 pp. <https://doi.org/10.2760/881843>
- Pumkao P, Takahashi J, Iwahashi H (2021) Detection and monitoring of insect traces in bioaerosols. *PeerJ* 9: e10862. <https://doi.org/10.7717/peerj.10862>
- Richardson RT, Lin CH, Quijia JO, Riusech NS, Goodell K, Johnson RM (2015) Rank-based characterization of pollen assemblages collected by honey bees using a multi-locus metabarcoding approach. *Applications in Plant Sciences* 3(11): 1500043. <https://doi.org/10.3732/apps.1500043>
- Richardson RT, Curtis HR, Matcham EG, Lin CH, Suresh S, Sponsler DB, Hearon LE, Johnson RM (2019) Quantitative multi-locus metabarcoding and waggle dance interpretation reveal honey bee spring foraging patterns in Midwest agroecosystems. *Molecular Ecology* 28(3): 686–697. <https://doi.org/10.1111/mec.14975>
- Ritter CD, Häggqvist S, Karlsson D, Sääksjärvi I, Muasya AM, Nilsson RH, Antonelli A (2019) Biodiversity assessments in the 21st century: The potential of insect traps to complement environmental samples for estimating eukaryotic and prokaryotic diversity using high-throughput DNA metabarcoding. *Genome* 62(3): 147–159. <https://doi.org/10.1139/gen-2018-0096>
- Roger F, Ghanavi HR, Danielsson N, Wahlberg N, Löndahl J, Pettersson LB, Andersson GKS, Boke Olén N, Clough Y (2022) Airborne environmental DNA metabarcoding for the monitoring of terrestrial insects – a proof of concept from the field. *Environmental DNA* 4(4): 790–807. <https://doi.org/10.1002/edn3.290>

- Ronquist F, Forshage M, Haggqvist S, Karlsson D, Hovmoller R, Bergsten J, Holston K, Britton T, Abenius J, Andersson B, Buhl PN, Coulianos CC, Fjellberg A, Gertsson CA, Hellqvist S, Jaschhof M, Kjaerandsen J, Klopstein S, Kobro S, Liston A, Meier R, Pollet M, Riedel M, Rohacek J, Schuppenhauer M, Stigenberg J, Struwe I, Taeger A, Ulefors SO, Varga O, Withers P, Gardenfors U (2020) Completing Linnaeus's inventory of the Swedish insect fauna: Only 5,000 species left? *PLoS ONE* 15(3): e0228561. <https://doi.org/10.1371/journal.pone.0228561>
- Schmidt A, Schneider C, Decker P, Hohberg K, Rombke J, Lehmitz R, Balint M (2022) Shotgun metagenomics of soil invertebrate communities reflects taxonomy, biomass, and reference genome properties. *Ecology and Evolution* 12(6): e8991. <https://doi.org/10.1002/ece3.8991>
- Schneider J, Valentini A, Dejean T, Montarsi F, Taberlet P, Glaizot O, Fumagalli L (2016) Detection of invasive mosquito vectors using environmental DNA (eDNA) from water samples. *PLoS ONE* 11(9): e0162493. <https://doi.org/10.1371/journal.pone.0162493>
- Seibold S, Gossner MM, Simons NK, Blüthgen N, Muller J, Ambarli D, Ammer C, Bauhus J, Fischer M, Habel JC, Linsenmair KE, Nauss T, Penone C, Prati D, Schall P, Schulze ED, Vogt J, Wollauer S, Weisser WW (2019) Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature* 574(7780): 671–674. <https://doi.org/10.1038/s41586-019-1684-3>
- Shirazi S, Meyer RS, Shapiro B (2021) Revisiting the effect of PCR replication and sequencing depth on biodiversity metrics in environmental DNA metabarcoding. *Ecology and Evolution* 11(22): 15766–15779. <https://doi.org/10.1002/ece3.8239>
- Shum P, Palumbi SR (2021) Testing small-scale ecological gradients and intraspecific differentiation for hundreds of kelp forest species using haplotypes from metabarcoding. *Molecular Ecology* 30(13): 3355–3373. <https://doi.org/10.1111/mec.15851>
- Simmons BI, Balmford A, Bladon AJ, Christie AP, De Palma A, Dicks LV, Gallego-Zamorano J, Johnston A, Martin PA, Purvis A, Rocha R, Wauchope HS, Wordley CFR, Worthington TA, Finch T (2019) Worldwide insect declines: An important message, but interpret with caution. *Ecology and Evolution* 9(7): 3678–3680. <https://doi.org/10.1002/ece3.5153>
- Sow A, Brévault T, Benoit L, Chapuis M, Galan M, Coeur d'acier A, Delvare G, Sembène M, Haran J (2019) Deciphering host-parasitoid interactions and parasitism rates of crop pests using DNA metabarcoding. *Scientific Reports* 9(1): 3646. <https://doi.org/10.1038/s41598-019-40243-z>
- Srivathsan A, Lee L, Katoh K, Hartop E, Kutty SN, Wong J, Yeo D, Meier R (2021) ONTbar-coder and MinION barcodes aid biodiversity discovery and identification by everyone, for everyone. *BMC Biology* 19(1): 217. <https://doi.org/10.1186/s12915-021-01141-x>
- Srivathsan A, Ang Y, Heraty JM, Hwang WS, Jusoh WFA, Kutty SN, Puniamoorthy J, Yeo D, Roslin T, Meier R (2023) Convergence of dominance and neglect in flying insect diversity. *Nature Ecology & Evolution* 7(7): 1012–1021. <https://doi.org/10.1038/s41559-023-02066-0>
- Swenson SJ, Gemeinholzer B (2021) Testing the effect of pollen exine rupture on metabarcoding with Illumina sequencing. *PLoS ONE* 16(2): e0245611. <https://doi.org/10.1371/journal.pone.0245611>
- Swift JF, Lance RF, Guan X, Britzke ER, Lindsay DL, Edwards CE (2018) Multifaceted DNA metabarcoding: Validation of a noninvasive, next-generation approach to studying bat populations. *Evolutionary Applications* 11(7): 1120–1138. <https://doi.org/10.1111/eva.12644>
- Tedersoo L, Drenkhan R, Anslan S, Morales-Rodriguez C, Cleary M (2019) High-throughput identification and diagnostics of pathogens and pests: Overview and practi-

- cal recommendations. *Molecular Ecology Resources* 19(1): 47–76. <https://doi.org/10.1111/1755-0998.12959>
- Theissinger K, Fernandes C, Formenti G, Bista I, Berg PR, Bleidorn C, Bombarely A, Crotini A, Gallo GR, Godoy JA, Jentoft S, Malukiewicz J, Mouton A, Oomen RA, Paez S, Palsboll PJ, Pampoulie C, Ruiz-Lopez MJ, Secomandi S, Svardal H, Theofanopoulou C, de Vries J, Waldvogel AM, Zhang G, Jarvis ED, Balint M, Ciofi C, Waterhouse RM, Mazzoni CJ, Hoglund J, Aghayan SA, Alioto TS, Almudi I, Alvarez N, Alves PC, Amorim do Rosario IR, Antunes A, Arribas P, Baldrian P, Bertorelle G, Böhne A, Bonisoli-Alquati A, Boštjančič LL, Boussau B, Breton CM, Buzan E, Campos PF, Carreras C, Castro LFIC, Chueca LJ, Čiampor F, Conti E, Cook-Deegan R, Croll D, Cunha MV, Delsuc F, Dennis AB, Dimitrov D, Faria R, Favre A, Fedrigo OD, Fernández R, Ficetola GF, Flot J-F, Gabalaldón T, Agius DR, Giani AM, Gilbert MTP, Grebenc T, Guschanski K, Guyot R, Hausdorf B, Hawlitschek O, Heintzman PD, Heinze B, Hiller M, Husemann M, Iannucci A, Irisarri I, Jakobsen KS, Klinga P, Kloch A, Kratochwil CF, Kusche H, Layton KKS, Leonard JA, Lerat E, Liti G, Manousaki T, Marques-Bonet T, Matos-Maraví P, Matschiner M, Maurmus F, Mc Cartney AM, Meiri S, Melo-Ferreira J, Mengual X, Monaghan MT, Montagna M, Mysłajek RW, Neiber MT, Nicolas V, Novo M, Ozretić P, Palero F, Pârvulescu L, Pascual M, Paulo OS, Pavlek M, Pegueroles C, Pellissier L, Pesole G, Primmer CR, Riesgo A, Rüber L, Rubolini D, Salvi D, Seehausen O, Seidel M, Studer B, Theodoridis S, Thines M, Urban L, Vasemägi A, Vella A, Vella N, Vernes SC, Vernesi C, Vieites DR, Wheat CW, Wörheide G, Wurm Y, Zammit G (2023) How genomics can help biodiversity conservation. *Trends in Genetics* 39(7): 545–559. <https://doi.org/10.1016/j.tig.2023.01.005>
- Thomsen PF, Sigsgaard EE (2019) Environmental DNA metabarcoding of wild flowers reveals diverse communities of terrestrial arthropods. *Ecology and Evolution* 9(4): 1665–1679. <https://doi.org/10.1002/ece3.4809>
- Tkacz A, Hortal M, Poole PS (2018) Absolute quantitation of microbiota abundance in environmental samples. *Microbiome* 6(1): 110. <https://doi.org/10.1186/s40168-018-0491-7>
- Uiterwaal SF, DeLong JP (2020) Using patterns in prey DNA digestion rates to quantify predator diets. *Molecular Ecology Resources* 20(6): 1723–1732. <https://doi.org/10.1111/1755-0998.13231>
- van der Heyde M, Bunce M, Wardell-Johnson G, Fernandes K, White NE, Nevill P (2020) Testing multiple substrates for terrestrial biodiversity monitoring using environmental DNA metabarcoding. *Molecular Ecology Resources* 00(3): 1–14. <https://doi.org/10.1111/1755-0998.13148>
- van Klink R, Bowler DE, Gongalsky KB, Swengel AB, Gentile A, Chase JM (2020) Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science* 368(6489): 417–420. <https://doi.org/10.1126/science.aax9931>
- van Klink R, August T, Bas Y, Bodesheim P, Bonn A, Fossey F, Hoye TT, Jongejans E, Menz MHM, Miraldo A, Roslin T, Roy HE, Ruczynski I, Schigel D, Schaffler L, Sheard JK, Svenningsen C, Tschan GF, Waldchen J, Zizka VMA, Astrom J, Bowler DE (2022a) Emerging technologies revolutionise insect ecology and monitoring. *Trends in Ecology & Evolution* 37(10): 872–885. <https://doi.org/10.1016/j.tree.2022.06.001>
- van Klink R, Bowler DE, Gongalsky KB, Chase JM (2022b) Long-term abundance trends of insect taxa are only weakly correlated. *Biology Letters* 18(2): 20210554. <https://doi.org/10.1098/rsbl.2021.0554>
- van Strien AJ, Termaat T, Groenendijk D, Mensing V, Kéry M (2010) Site-occupancy models may offer new opportunities for dragonfly monitoring based on daily species lists. *Basic and Applied Ecology* 11(6): 495–503. <https://doi.org/10.1016/j.baae.2010.05.003>

- van Strien AJ, van Swaay CAM, Termaat T, Devictor V (2013) Opportunistic citizen science data of animal species produce reliable estimates of distribution trends if analysed with occupancy models. *Journal of Applied Ecology* 50(6): 1450–1458. <https://doi.org/10.1111/1365-2664.12158>
- Vasselon V, Ács É, Almeida S, Andree K, Apothélos-Perret-Gentil L, Bailet B, Baricevic A, Beentjes K, Bettig J, Bouchez A, Capelli C, Chardon C, Duleba M, Elersek T, Genthon C, Hurtz M, Jacas L, Kahlert M, Kelly M, Lewis M, Macher J, Mauri F, Moletta-Denat M, Mortágua A, Pawlowski J, Pérez Burillo J, Pfannkuchen M, Pilgrim E, Pissaridou P, Porter J, Rimet F, Stanic K, Tapolczai K, Theroux S, Trobajo R, van der Hoorn B, Vasquez Hadjilyra M, Walsh K, Wanless D, Warren J, Zimmermann J, Zupančič M (2021) The Fellowship of the Ring Test: DNAqua-Net WG2 initiative to compare diatom metabarcoding protocols used in routine freshwater biomonitoring for standardisation. *ARPHA Conference Abstracts 4*: e65142. <https://doi.org/10.3897/aca.4.e65142>
- Wang J, Santiago E, Caballero A (2016) Prediction and estimation of effective population size. *Heredity* 117(4): 193–206. <https://doi.org/10.1038/hdy.2016.43>
- Wang WY, Srivathsan A, Foo M, Yamane SK, Meier R (2018) Sorting specimen-rich invertebrate samples with cost-effective NGS barcodes: Validating a reverse workflow for specimen processing. *Molecular Ecology Resources* 18(3): 490–501. <https://doi.org/10.1111/1755-0998.12751>
- Watts C, Dopheide A, Holdaway R, Davis C, Wood J, Thornburrow D, Dickie IA (2019) DNA metabarcoding as a tool for invertebrate community monitoring: A case study comparison with conventional techniques. *Austral Entomology* 58(3): 675–686. <https://doi.org/10.1111/aen.12384>
- Weitemier K, Penaluna BE, Hauck LL, Longway LJ, Garcia T, Cronn R (2021) Estimating the genetic diversity of Pacific salmon and trout using multigene eDNA metabarcoding. *Molecular Ecology* 30(20): 4970–4990. <https://doi.org/10.1111/mec.15811>
- Wilson RS, Keller A, Shapcott A, Leonhardt SD, Sickel W, Hardwick JL, Heard TA, Kaluza BF, Wallace HM (2021) Many small rather than few large sources identified in long-term bee pollen diets in agroecosystems. *Agriculture, Ecosystems & Environment* 310: 107296. <https://doi.org/10.1016/j.agee.2020.107296>
- Wühl L, Pylatiuk C, Giersch M, Lapp F, von Rintelen T, Balke M, Schmidt S, Cerretti P, Meier R (2022) DiversityScanner: Robotic handling of small invertebrates with machine learning methods. *Molecular Ecology Resources* 22(4): 1626–1638. <https://doi.org/10.1111/1755-0998.13567>
- Yang C, Bohmann K, Wang X, Cai W, Wales N, Ding Z, Gopalakrishnan S, Yu DW (2020) Biodiversity Soup II: A bulk-sample metabarcoding pipeline emphasizing error reduction. *BioRxiv*. <https://doi.org/10.1101/2020.07.07.187666>
- Young RG, Milian-Garcia Y, Yu J, Bullas-Appleton E, Hanner RH (2021) Biosurveillance for invasive insect pest species using an environmental DNA metabarcoding approach and a high salt trap collection fluid. *Ecology and Evolution* 11(4): 1558–1569. <https://doi.org/10.1002/ece3.7113>
- Yu DW, Ji Y, Emerson BC, Wang X, Ye C, Yang C, Ding Z (2012) Biodiversity soup: Metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods in Ecology and Evolution* 3(4): 613–623. <https://doi.org/10.1111/j.2041-210X.2012.00198.x>
- Zenker MM, Specht A, Fonseca VG (2020) Assessing insect biodiversity with automatic light traps in Brazil: Pearls and pitfalls of metabarcoding samples in preservative ethanol. *Ecology and Evolution* 10(5): 2352–2366. <https://doi.org/10.1002/ece3.6042>
- Zinger L, Bonin A, Alsos IG, Balint M, Bik H, Boyer F, Chariton AA, Creer S, Coissac E, Deagle BE, De Barba M, Dickie IA, Dumbrell AJ, Ficetola GF, Fierer N, Fumagalli L,

- Gilbert MTP, Jarman S, Jumpponen A, Kauserud H, Orlando L, Pansu J, Pawlowski J, Tedersoo L, Thomsen PF, Willerslev E, Taberlet P (2019) DNA metabarcoding-Need for robust experimental designs to draw sound ecological conclusions. *Molecular Ecology* 28(8): 1857–1862. <https://doi.org/10.1111/mec.15060>
- Zinger L, Donald J, Brosse S, Gonzalez MA, Iribar A, Leroy C, Muriene J, Orivel J, Schimann H, Taberlet P, Lopes CM (2020) Advances and prospects of environmental DNA in neotropical rainforests. In: Dumbrell AJ, Turner EC, Fayle TM (Eds) *Advances in ecological research*, 331–373. <https://doi.org/10.1016/bs.aecr.2020.01.001>
- Zizka VMA, Leese F, Peinert B, Geiger MF (2019) DNA metabarcoding from sample fixative as a quick and voucher-preserving biodiversity assessment method (1). *Genome* 62(3): 122–136. <https://doi.org/10.1139/gen-2018-0048>

Supplementary material 1

Background information

Authors: Wiebke Sickel, Vera Zizka, Alice Scherges, Sarah J. Bourlat, Petra Dieker

Data type: docx

Explanation note: Background information on DNA metabarcoding and methodological adjustments to improve abundance and biomass estimates.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mbmg.7.112290.suppl1>

Supplementary material 2

Literature collection

Authors: Wiebke Sickel, Vera Zizka, Alice Scherges, Sarah J. Bourlat, Petra Dieker

Data type: xlsx

Explanation note: Relevant publications were identified in two steps: 1) via an online literature search in Google Scholar and EBSCO Discovery Service; keywords: ((quant*) AND (insect) AND (metabarcod*) AND (DNA)), including only peer-reviewed publications in English, results were screened for suitability based on title and abstract; 2) addition of publications based on the authors' expertise; included studies that applied metabarcoding with arthropods and/or in relation to their trophic interactions within ecosystems (e.g. pollination, food web studies); excluded topics were: individual-based DNA barcoding and NGS barcoding, long-read sequencing methodology, mito-/metagenomics and genome skimming.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mbmg.7.112290.suppl2>

Supplementary material 3

Categories assessed in the literature review

Authors: Wiebke Sickel, Vera Zizka, Alice Scherges, Sarah J. Bourlat, Petra Dieker

Data type: xlsx

Explanation note: For each category, possible parameters are given, together with examples and more detailed explanation where appropriate and necessary.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mbmg.7.112290.suppl3>

Supplementary material 4

Evaluation of methodological approaches

Authors: Wiebke Sickel, Vera Zizka, Alice Scherges, Sarah J. Bourlat, Petra Dieker

Data type: xlsx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mbmg.7.112290.suppl4>